

MANAGEMENT PROTOCOL OF CHILDREN WITH COVID-19

Approved Date: 2078/09/14

Approved by Hon. Minister, Ministry of Health and Population



Government of Nepal
Ministry of Health and Population

FOREWORD



Ref. No.

Government of Nepal
Ministry of Health
DEPARTMENT OF HEALTH SERVICES



Tel. : 4261436
: 4261712
Fax : 4262238

Pachali, Teku
Kathmandu, Nepal

Date:-.....

Forewords

With immense pleasure I am writing few words of appreciation on the timely development of "Management protocol of children with COVID-19" led by Curative Service Division. Various variants have been identified and have been classified as Variants of Interest or Variants of Concern. The recently identified Variant of Concern, Omicron is a highly divergent variant with a high number of mutations, which has uncertainties regarding its transmissibility and severity. Due to a smaller number of children being immunized, they are still at higher risk of infection. A guidance document for proper management of COVID-19 among pediatric population was lacking, which has been fulfilled by the timely development of "Management protocol of children with COVID-19". This management protocol is expected to help relevant healthcare workers to properly manage the pediatric cases and to standardize the management of pediatric cases.

I would like to express my sincere gratitude to Curative Service Division for taking the lead and the experts of Nepal Pediatric Society (NEPAS) including all the contributors for their tireless effort on bringing forth "Management protocol of children with COVID-19". I am grateful to WHO and UNICEF for their technical and financial support in development of this concise yet comprehensive management protocol.


.....

Dr. Dipendra Raman Singh

Director General

Department of Health Services

FOREWORD



Ref. No.

Government of Nepal
Ministry of Health
DEPARTMENT OF HEALTH SERVICES
(.....)
Curative Service Division
Kathmandu

Tel. : 4261436
: 4261712
Fax : 4262238

Pachali, Teku
Kathmandu, Nepal

Date:-.....

Foreword

I feel privileged to put some words on "Management protocol of children with COVID-19". The emergence of various variants with increased transmissibility and disease severity, countries in the world have experienced several waves with increased hospitalization and deaths. In addition to the widely circulating Delta Variant, the emergence of new variant Omicron with a high number of mutations is quite concerning. The likelihood of potential introduction and subsequent transmission of Omicron in the South-East Asia region is high. Children are still at higher risk of infection as most of them are still non-immunized. Development of management protocol helps to bring uniformity in management of cases. This management protocol is expected to provide guidance to relevant healthcare workers for proper management of the pediatric cases.

I would like to express my sincere gratitude to the Section Chiefs, all the staffs of Curative Service Division, the experts of Nepal Pediatric Society (NEPAS) and the contributors including WHO and UNICEF for their technical and financial support in development of this comprehensive management protocol.

Dr. Pawan Jung Rayamajhi

Director

Curative Service Division, DoHS

CORE WORKING TEAM

Dr. Jyoti Ratna Dhakwa
Senior Consultant Pediatrician,
Ishan Children and women's Hospital

Dr. Krishna Prasad Poudel
Director EDCD, Teku

Dr. Krishna Prasad Bista
Consultant Pediatrician,
Kanti Children's Hospital

Dr. Nisha Jyoti Shrestha
Chief Consultant Pediatrician
Kanti Children's Hospital
Assistant Professor, Department of
Pediatrics, NAMS

Dr. Ram Hari Chapagain
Senior consultant pediatrician,
Kanti Children's Hospital
Associate Professor, Department of
Pediatrics, NAMS

Dr. Puja Amatya
Pediatric Intensive Care Specialist
Assistant Professor, Patan Academy of
Health Sciences, Lalitpur

Dr. Sumit Agrawal
Senior Consultant Pediatrician
Kanti Children's Hospital

Dr. Pradyuman Prasad Chauhan
Medical Officer, HEOC
Ministry of Health and Population

Dr. Netra Rana
Consultant Pediatrician
Lumbini Hospital

Dr. Biraj Parajuli
Pediatric Intensivist
Assistant Professor, Chitwan Medical College

Dr. Ganesh Kumar Rai
President, NEPAS

Dr. Arun Kumar Sharma
Associate Professor,
Institute of Medicine, Tribhuvan University

Dr. Moon Thapa
Associate professor & HOD,
Department of Pediatrics, Army Hospital

Dr. Yuba Nidhi Basaula
Senior Consultant Pediatrician
Bharatpur Hospital

Dr. Mukesh Bhatta
Assistant Professor, Pediatrics,
BPKIHS, Dharan

Dr. Narmada Devkota (M.Phil., PhD)
Clinical Psychologist
Child and Adolescent Psychiatry Unit,
Kanti Children's Hospital

Dr. Santosh Adhikari
Consultant Pediatrician,
Kanti Children's Hospital

Dr. Rupesh Shrestha
Pediatrician,
TUTH

Dr. Raju Kafle
Associate Professor,
UCMS

Dr. Roma Bora
Associate Professor,
Nepalgunj Medical College

Dr Samana Sharma
Consultat paediatrician,
Scheer Memorial Hospital, Banepa

TABLE OF CONTENTS

1. Infection Prevention and Control (IPC) for COVID-19.....	1
2. Suspicion, Testing and Management of COVID-19 in Children.....	16
3. Neonates with COVID-19.....	34
4. Multi-System Inflammatory Syndrome in Children (MIS-C).....	39
5. Crisis Responses and Psycho-social Management for Children and Adolescents (C&A) during COVID-19 Pandemic.....	46
6. Managing Children with Disabilities in COVID-19 Facilities: Guidance for Doctors, Nurses, Paramedic and other Health Care Staff.....	48
7. Integrating Gender, Equity and Human Rights (GER) in Clinical Management (CM) of COVID-19.....	52
Appendix 1. Figures.....	57
Appendix 2. Flow charts.....	65
Appendix 3. Multisystem Inflammatory Syndrome in Children (MIS-C).....	70
References.....	74
List of Contributors.....	79

CHAPTER I

INFECTION PREVENTION AND CONTROL (IPC) FOR COVID-19

1. Background

Infection prevention and control (IPC), an essential part of health care infrastructure, is the practical discipline concerned with preventing health-care-associated infection.

The five IPC strategies required to prevent or limit transmission of COVID-19 in health care facilities include:

1. Screening and triage for early recognition of patients with suspected COVID-19 and rapid implementation of source control measures
2. Applying standard precautions for all patients
3. Implementing additional precautions
4. Implementing administrative controls
5. Implementing environmental and engineering controls

1.1. Screening and Triage for Early Recognition of Patients with Suspected COVID-19 and Rapid Implementation of Source Control Measures

To facilitate screening and triage, health-care facilities should:

Establish entrances and display information for patients with signs and symptoms of COVID-19 to report to designated area for screening.

Train staffs on the signs and symptoms of COVID-19 and the most recent case definitions.

Encourage health care workers (HCWs) to be alert to potential COVID-19 infection in all patients.

Establish well-equipped screening and triage stations.

Ensure that screening personnel maintain a distance of at least 2 meter from patients, ideally with a separation created by a glass/plastic screen. If not possible, mask and eye protection should be worn.

Use a screening algorithm to promptly identify and direct patients with suspected COVID-19 to an isolation room or dedicated COVID-19 waiting area; all suspected COVID-19 patients should wear masks for source control purposes and be positioned at least 2 meter apart from each other in a designated, well-ventilated, waiting area.

For Isolation or designated waiting area:

Health-care facilities without enough single isolation rooms in emergency departments should designate a separate, well-ventilated area (with benches/chairs placed at least 2 meter apart) where suspected COVID-19 patients can wait.

The area should have toilets, hand hygiene stations, and trash bins with lid for disposal of paper tissues and display graphic information on how to perform hand and respiratory hygiene.

1.2. Applying standard precautions for all patients

Standard precautions represent the basic level of infection control measures that should be used at all times in the care of all patients. Standard precautions include, but are not limited to, hand and respiratory hygiene, the use of appropriate personal protective equipment (PPE) according to risk assessment, environmental cleaning, and safe waste management.

1.2.1. Hand hygiene is one of the most effective measures to prevent the spread of COVID-19 and other pathogens. For optimal hand hygiene performance, following principles should be applied:

- Perform hand hygiene according to the WHO's My 5 Moments for Hand Hygiene (Figure 1) approach in the following five situations: before touching a patient, before performing any clean or aseptic procedure, after exposure to body fluid, after touching a patient, and after touching a patient's surroundings.
- Hand hygiene includes either cleansing hands with an alcohol-based hand rub (ABHR) containing at least 70% alcohol (for 20-30 seconds), or with soap, water and disposable towels (for 40-60 seconds).

- | |
|--|
| <ul style="list-style-type: none"> • Alcohol-based hand rub products are preferred if hands are not visibly soiled. |
| <ul style="list-style-type: none"> • Wash hands with soap and water (Figure 2) when they are visibly soiled. |

1.2.2 Respiratory hygiene (Figure 3) measures to be ensured are:

- | |
|---|
| <ul style="list-style-type: none"> • Display graphic information on the need to cover nose and mouth with a tissue or bent elbow when coughing or sneezing. |
| <ul style="list-style-type: none"> • Perform hand hygiene after contact with respiratory secretions or objects that may be potentially contaminated with respiratory secretions. |
| <ul style="list-style-type: none"> • Give patients with suspected COVID-19 a medical mask to wear. |

Children aged up to five years should not wear masks for source control. The rationale includes consideration of the fact that by the age of five years, children usually achieve significant developmental milestones, including the manual dexterity and fine motor coordination movements needed to appropriately use a mask with minimal assistance.

1.2.3. Use of personal protective equipment (PPE):

The rational and correct use of PPE (Table 1) reduces exposure to pathogens. The effectiveness of PPE strongly depends on:

- Staff training on putting on and removing PPE;
- Prompt access to sufficient supplies;
- Appropriate hand hygiene;
- Health worker compliance;
- Regular monitoring and feedback by IPC personnel.

Steps to put on (don) and remove (doff) PPE are included in Figure 4 and 5 of Appendices.

Table 1. Recommended type of PPE to be used in the context of COVID-19 disease, according to the setting, personnel and type of activity ^a

Targeted personnel	Activity	Medical Mask	Gown	Gloves	Eye Protection	Respirator (N95/ FFP2)	Apron	Boots	Remarks
Healthcare facilities (Inpatient facilities): Patient room									
Health-care workers	Providing direct care to COVID-19 patients	✓	✓	✓	✓	✗	✗	✗	
	Aerosol-generating procedures	✓	✓	✓	✓	✓	✓		
Cleaners	Entering the room of COVID-19 patients	✓	✓	✓ (heavy duty)	✓	✗	✗	✓	eye protection if risk of splash from organic material or chemicals
Visitors ^b	Entering the room of a COVID-19 patient	✓	✓	✓	✗	✗	✗	✗	
Inpatient facilities: Other areas of patient transit (e.g., wards, corridors)									
All staffs & HCWs	Any activity that does not involve contact with COVID-19 patients	✓	✗	✗	✗	✗	✗	✗	Universal masking in areas of known or suspected community or cluster SARS-CoV-2 transmission & Targeted continuous masking in areas where SARS-CoV-2 transmission is known or suspected to be only sporadic
Inpatient and Outpatient facilities: Triage									
HCWs	Preliminary screening not involving direct contact ^c	✓	✗	✗	✗	✗	✗	✗	Universal masking in areas of known or suspected community or cluster
Patients with respiratory symptoms	Any	✓	✗	✗	✗	✗	✗	✗	SARS-CoV-2 transmission & Targeted continuous masking in areas where SARS-CoV-2 transmission is known or suspected to be only sporadic
Patients without respiratory symptoms	Any	✓	✗	✗	✗	✗	✗	✗	

Inpatient facilities: Laboratory									
Lab Technician	Manipulation of respiratory samples	✓	✓	✓	✓	✗	✗	✗	Eye protection if risk of splash
Healthcare facilities (Outpatient facilities): Consultation room									
Healthcare workers	Examination of patients with respiratory symptoms	✓	✓	✓	✓	✗	✗	✗	
HCWs	Examination of patients without respiratory symptoms	✓							PPE as per standard precautions & risk assessment
Patients with respiratory symptoms	Any	✓							Provide medical mask if tolerated
Patients without respiratory symptoms	Any	✓							Universal masking in areas of known or suspected community or cluster SARS-CoV-2 transmission
Cleaners	After and between consultations with patients with respiratory symptoms	✓	✓	✓ (heavy duty)	✓			✓	Eye protection if splash risk from organic material or chemicals
Outpatient facilities: Waiting room									
Patients with respiratory symptoms	Any	Provide medical mask if tolerated; Immediately move the patient to an isolation room; if not feasible, ensure spatial distance of at least 2 meter from other patients							
Patients without respiratory symptoms	Any	✓							Universal masking in areas of known or suspected community or cluster SARS-CoV-2 transmission
Administrative areas									
All staff, including HCWs	Administrative tasks	✓							Universal masking in areas of known or suspected community or cluster SARS-CoV-2 transmission

Management protocol of Children with COVID-19

Ambulance or transfer vehicle									
Healthcare workers	Transporting suspected COVID-19 patients to the referral healthcare facility	✓	✓	✓	✓				
Driver	Involved only in driving the suspected patient and the compartment separated	✓							Universal masking in areas of known or suspected community or cluster SARS-CoV-2 transmission
	Assisting with loading/unloading of suspected patient	✓	✓	✓	✓				
	No direct contact with patient but no separation between compartment	✓							
Patient with suspected COVID-19	Transport to the referral healthcare facility	✓							Medical mask if tolerated
Cleaners	Cleaning after and between transport of patients with suspected COVID-19 disease to the referral healthcare facility	✓	✓	✓	✓			✓	

^aIn addition to using appropriate PPE, frequent hand hygiene & respiratory hygiene should always be performed. PPE should be discarded in an appropriate waste container after use, and hand hygiene performed before putting on (donning) and after taking off PPE (doffing).

^bNumber of visitors should be restricted. If visitors must enter a COVID-19 patient's room, they should wear gown, mask and perform hand hygiene.

^cThis category includes the use of no-touch thermometers, thermal imaging cameras, and limited observation and questioning, all while maintaining a spatial distance of at least 2m.

Universal masking in health facilities is defined as the requirement for all persons (staff, patients, visitors, service providers and others) to wear a mask at all times except for when eating or drinking. All health workers, including community health workers and caregivers, should wear a medical mask at all times, for any activity (care of COVID-19 or non-COVID-19 patients) and in any common area (e.g., cafeteria, staff rooms). Other staffs, visitors, outpatients and service providers should also wear a mask (medical or non-medical) at all times when in the health facility.

Targeted continuous medical mask use is defined as the practice of wearing a medical mask by all health workers and caregivers working in clinical areas during all routine activities throughout the entire shift. Health workers, including community health workers and caregivers who work in clinical areas, have to continuously wear a medical mask during routine activities throughout the entire shift, apart from when eating and drinking and changing their medical masks after caring for a patient who requires droplet/contact or airborne precautions for reasons other than COVID-19. In these transmission settings, staff working in non-patient areas (e.g. administrative staff) are not required to wear a medical mask during routine activities if they have no patient contact.

1.2.4. Environmental cleaning (Table 2)

- Clean surfaces thoroughly with water and detergent;
- Apply a disinfectant solution, either 0.1% (1000ppm) sodium hypochlorite or 70-90% ethanol. However, if there are large spills of blood or body fluids, a concentration of 0.5% (5000ppm) sodium hypochlorite should be used;
- Contact time of a minimum of 2 minute is recommended for ethanol, chlorine-based products and hydrogen peroxide $\geq 0.5\%$;
- After appropriate contact time, disinfectant residue may be rinsed off with clean water if required.
- Medical devices and equipment, laundry, food service utensils and medical waste should be managed in accordance with safe routine procedures.

Table 2. Health-care setting: Recommended frequency of cleaning of environmental surfaces, according to the patient areas with suspected or confirmed COVID-19 patients.

Patient area	Frequency ^a	Additional guidance
Screening/triage area	At least twice daily	Focus on high-touch surfaces, then floors (last).
Inpatient rooms / cohort- occupied	At least twice daily, preferably three times daily, in particular for high-touch surfaces	Focus on high-touch surfaces, starting with shared/common surfaces, then move to each patient bed; use new cloth for each bed if possible; then floors (last).
Inpatient rooms- unoccupied (terminal cleaning)	Upon discharge/ transfer	Low-touch surfaces, high-touch surfaces, floors (in that order); waste and linens removed, bed thoroughly cleaned and disinfected.
Outpatient / ambulatory care rooms	After each patient visit (in particular for high-touch surfaces) and at least once daily terminal clean	High-touch surfaces to be disinfected after each patient visit. Once daily low-touch surfaces, high-touch surfaces, floors (in that order); waste and linens removed, examination bed thoroughly cleaned and disinfected.

Hallways/corridors	At least twice daily ^b	High-touched surfaces including railings and equipment in hallways, then floors (last).
Patient bathrooms/ toilets	Private patient room toilet: at least twice daily Shared toilets: at least three times daily	High-touch surfaces, including door handles, light switches, counters, faucets, sink bowls, then toilets and finally floor (in that order). Avoid sharing toilets between staff and patients.

^aEnvironmental surfaces should also be cleaned and disinfected whenever visibly soiled or if contaminated by body fluid (e.g., blood); ^bFrequency can be once a day if hallways are not frequently used.

1.2.5. Waste Management

Health-care waste produced during the care of patients with suspected or confirmed COVID-19 is considered to be infectious and should be collected safely in clearly marked lined containers and sharp safe boxes.

To safely manage health-care waste, facilities should:

- Assign responsibility and adequate human and material resources to segregate and dispose of waste.
- Treat waste preferably on-site, and then safely dispose them. If waste is moved off-site, it is critical to understand where and how it will be treated and disposed.
- Use appropriate PPE (boots, long-sleeved gown, heavy-duty gloves, mask, and goggles or a face shield) while managing infectious waste and perform hand hygiene after doffing.
- Prepare for increases in the volume of infectious waste during the COVID-19 outbreak, especially through the use of PPE.

1.3. Implementing additional precautions

Transmission of the COVID-19 virus may occur by direct contact with infected people (respiratory droplets and contact routes) and indirect contact with surfaces in the immediate environment or with objects used on the infected person (e.g. stethoscope or thermometer). Airborne transmission of the COVID-19 virus is possible under circumstances and settings where aerosol generating procedures (AGPs) are performed.

Apart from standard precautions, additional precautions that can be implemented for COVID-19:

Isolation and cohorting of patients with suspected or confirmed COVID-19 in single rooms or, if unavailable, cohorting them in the same room with the following principles:

Designate a team of health workers, where possible, for care of patients with suspected or confirmed COVID-19 to reduce the risk of transmission.

Restrict the number of health workers in contact with each COVID-19 patient.

Patients should be placed in well ventilated single rooms if feasible.

When single rooms are not available, suspected, probable or confirmed COVID-19 patients should be grouped together (cohorted) in adequately ventilated areas with beds placed at least 2 meter apart (e.g. suspected with suspected).

Avoid moving and transporting patients out of their room or area unless medically necessary. Use designated portable X-ray equipment and/or other designated diagnostic equipment.

If transport is required, use predetermined transport routes to minimize exposure for staff, other patients and visitors, and give the patient a medical mask to wear if tolerated.

Ensure that HCWs who are transporting patients perform hand hygiene and wear appropriate PPE.

Equipment should be either single-use and disposable or dedicated equipment (e.g. stethoscopes, blood pressure cuffs and thermometers). If equipment needs to be shared between patients, clean and disinfect after each use.

Maintain a record of all staff entering the patient's room.

Contact and droplet precautions in addition to standard precautions should be followed, with the following principles:

Perform hand hygiene before donning and after doffing; use appropriate PPE.

HCWs and caregivers working in clinical areas in areas with community transmission COVID-19 should continuously wear a medical mask during all routine activities throughout the entire shift.

It is not necessary for HCWs and caregivers to wear boots, coverall and apron during routine care.

Extended use of medical mask, gown and eye protection can be applied during the care of COVID-19 patients during PPE shortages.

For a COVID-19 patient who is infected with a multi-drug resistant organism (e.g. *Clostridioides difficile*), a new set of gown and gloves are needed after caring for such patients.

HCWs should refrain from touching their eyes, nose or mouth with potentially contaminated gloved or bare hands.

Notify the area receiving the patient of any necessary precautions before the patient's arrival.

Frequently clean and disinfect surfaces with which the patient is in contact.

Some AGPs have been associated with an increased risk of transmission of coronaviruses (SARS-CoV-1, SARS-CoV-2 and MERS-CoV). The current WHO list of these AGPs is: tracheal intubation, non-invasive ventilation (e.g. BiPAP, CPAP), tracheotomy, cardiopulmonary resuscitation, manual ventilation before intubation, bronchoscopy, sputum induction induced by using nebulized hypertonic saline, and autopsy procedures. It remains unclear whether aerosols generated by nebulizer therapy or high-flow oxygen delivery are infectious.

Perform procedures in an adequately ventilated room.

Use appropriate PPE: wear a particulate respirator at least as protective as a US National Institute for Occupational Safety and Health (NIOSH)-certified N95, European Union (EU) standard FFP2, or equivalent.

Although initial fit testing is needed prior to the use of a particulate respirator, many countries and health-care facilities do not have a respiratory fit testing programme. Therefore, it is critical that when health workers put on a disposable particulate respirator, they should always perform the required seal check to ensure there is no leakage.

Other PPE items include eye protection (i.e. goggles or a face shield), long-sleeved gown and gloves. If gowns are not fluid resistant, HCWs performing AGPs should use a waterproof apron if the procedure is expected to produce a large volume of fluid that might penetrate the gown.

In the intensive care units, where AGPs are frequently performed, the HCW may choose to wear a particulate respirator throughout his or her shift, in areas of community transmission; keep the number of persons present in the room to the absolute minimum required for the patient's care.

1.4. Implementing administrative controls:

Administrative controls and policies for the prevention and control of transmission of COVID-19 within the health-care facility includes:

- Establishing sustainable IPC infrastructures and activities.
- Educating patients' caregivers.
- Developing policies for early recognition of patients with suspected COVID-19.
- Ensuring access to laboratory testing for COVID-19 detection.
- Preventing overcrowding, especially in the emergency department.
- Providing dedicated waiting areas for symptomatic patients.
- Planning for (e.g. repurposing of other wards) and isolating COVID-19 patients.
- Ensuring adequate supplies of PPE.
- Ensuring adherence to IPC policies and procedures in all aspects of health care.

1.5. Implementing environmental and engineering controls

Environmental and engineering controls, an integral part of IPC, aims to reduce the concentration of infectious respiratory aerosols (i.e. droplet nuclei) in the air and the contamination of surfaces and inanimate objects. They include standards for adequate ventilation according to specific areas in health-care facilities, adapted structural design, spatial separation, as well as adequate environmental cleaning.

Important Considerations

A. Collecting and handling laboratory specimens from patients with suspected COVID-19:

All specimens collected for laboratory investigations should be regarded as potentially infectious. Health workers who collect, handle or transport any clinical specimens should adhere to the following measures and biosafety practices to minimize the possibility of exposure to pathogens:

Ensure that HCWs who collect specimens, including nasopharyngeal and oropharyngeal swabs, use appropriate PPE (i.e. eye protection, a medical mask, a long-sleeved gown and gloves). If the specimen is collected with an AGP (e.g. sputum induction), personnel conducting the procedure should wear a particulate respirator at least as protective as a NIOSH-certified N95, an EU standard FFP2, or equivalent.

Ensure that all personnel who transport specimens are trained in safe handling practices and spill decontamination procedures.

Place specimens for transport in leak-proof specimen bags (i.e. secondary containers) that have a separate sealable pocket for the specimen (i.e. a plastic biohazard specimen bag), with the patient's label on the specimen container (i.e. the primary container), and a clearly written laboratory request form.

Ensure that laboratories adhere to appropriate biosafety practices and transport requirements based on WHO's interim Laboratory biosafety guidance related to COVID-19.

Deliver all specimens by hand whenever possible.
Do not use pneumatic-tube systems to transport specimens.

Document clearly patient's full name, date of birth and clinical diagnosis of the suspected case of COVID-19 on the laboratory request form.
Notify the relevant laboratory as soon as possible that the specimen is being transported.

B. Considerations for surgical procedures:

In the context of the COVID-19 pandemic, every surgical procedure may entail risk for both health workers and patients. Any decision on whether to operate on a patient should not be based on the patient's COVID-19 status, but on need (e.g. trauma or emergency), the risks and benefits of surgery. The patient's clinical condition may deteriorate and develop life-threatening complications and outcomes if surgery is delayed.

Recent data point to a high proportion of post-operative pulmonary complications associated with increased mortality in patients with COVID-19.

C. IPC measures recommended for outpatient care:

<ul style="list-style-type: none">• Apply the basic principles of IPC and standard precautions in all health-care facilities, including outpatient settings and primary care.
<ul style="list-style-type: none">• Consider alternatives to face-to-face outpatient visits using telemedicine (e.g. telephone consultations or cell phone videoconference) to provide clinical support without direct contact with the patient.
<ul style="list-style-type: none">• Screening, early recognition and isolation of patients with suspected COVID-19.
<ul style="list-style-type: none">• Emphasis on hand hygiene, respiratory hygiene and medical masks to be used by patients with respiratory symptoms.
<ul style="list-style-type: none">• Appropriate use of contact and droplet precautions when performing clinical examination on patients with suspected COVID-19.
<ul style="list-style-type: none">• When symptomatic patients are required to wait, ensure they have a separate waiting area where patients can sit at least 2 meter apart and provide them with masks.

D. Dead body management:

Health workers should do a preliminary evaluation and risk assessment before undertaking any activity related to the management of suspected or confirmed COVID-19 fatality and follow WHO's IPC guidance for safe management of dead bodies in the context of COVID-19.

Health workers should:

- Perform hand hygiene before and after handling the body.
- Use appropriate PPE based on the level of interaction with the body and risk assessment (e.g. use of eye protection and medical masks in addition to gloves and fluid-resistant gown or apron, if there is a risk of body fluids splashes while handling the body).
- Ensure that any body fluids leaking from orifices are contained and cover body in cloth to transfer to mortuary area
- Not be engaged in any other activity during body handling or preparation.
- Disinfect any non-disposable equipment used during handling of the body as per WHO guidance.
- Correctly remove and dispose of PPE when finished.

Body bags are not necessary for COVID-19, although they may be used for other reasons such as excessive body fluid leakage or absence of refrigerated morgue, especially in countries with a warm climate. If more than 24 hours has passed since the person died, or if burial/cremation is not foreseen within the next 24-48 hours, a second body bag may be used.

CHAPTER II

SUSPICION, TESTING AND MANAGEMENT OF COVID-19 IN CHILDREN

2.1. When to suspect COVID-19 infection in children?

If the **children** are having signs/symptoms like:

Most common symptoms:

- Fever
- Dry cough
- Tiredness

Less common symptoms:

- Aches and pains
- Sore throat
- Diarrhea
- Conjunctivitis
- Headache
- Loss of taste or smell
- A rash on skin, or discoloration of fingers or toes

Serious symptoms:

- Difficult breathing or shortness of breath
- Chest pain or pressure
- Loss of speech or movement

Infants may have following additional symptoms:

- Diarrhea
- Vomiting
- Decreased feeding

2.2. Diagnosis of suspected children with COVID-19 symptoms

Indications for testing:

- A child with acute respiratory illness (fever and cough or fever and shortness of breath) AND new loss of smell OR taste.
- A child with acute respiratory illness (fever and cough or fever and shortness of breath) AND any two of these (chills, muscle pain, diarrhea, sore throat).
- A child with acute respiratory illness (fever and cough or fever and shortness of breath) in the absence of an alternative diagnosis that fully explains the clinical presentation.
- **Fever (>3 days) AND two of the following:**
 - a. Rash, non-purulent conjunctivitis or muco-cutaneous inflammation;
 - b. Hypotension or shock;
 - c. New cardiac abnormalities;
 - d. New bleeding disorder; and
 - e. Diarrhea, vomiting or abdominal pain.
- A child with acute respiratory illness (fever and cough or fever and shortness of breath) with underlying chronic conditions, immunocompromised conditions.
- Presentation with severe illness (e.g. new requirement for supplemental oxygen or increased requirement from baseline, new or increased need for ventilation [invasive or noninvasive] or clinical manifestations of multisystem inflammatory syndrome in children.
- **Children who were in close contact** (contact of at least 15 minutes over a 24-hour period within a distance of less than 6 feet) **with a person with confirmed or probable SARS-CoV-2 infection, and developed ANY of the symptoms described above.**
- Children who require screening, i.e. prior to a medical procedure such as elective surgery or as a school or workplace requirement.
- Infant born to a mother with suspected or confirmed COVID-19.
- If the treating clinician suspects COVID-19.

2.2.1. Testing Modalities

a. Reverse-Transcription Polymerase Chain Reaction (RT-PCR):

It is the gold standard for the diagnosis of COVID-19 in children. It is performed using nasopharyngeal swab specimen. However, false negative may be present. In the presence of strong clinical suspicion but negative initial PCR test, a repeat test is warranted to rule out COVID-19. The repeat testing should be performed after 24-48 hour of the first test.

b. Antigen Testing for SARS-CoV-2

Antigen tests are performed on nasopharyngeal swab specimens placed directly into the assay's extraction buffer or reagent. Antigen tests can be used as a point of care test and are inexpensive as compared to RT-PCR.

c. Antibody testing

It is useful for diagnosis of prior infection (or infection of at least 3 to 4 weeks' duration). The specimen used is blood.

Possible Indications of Antibody testing

1. In case of suspected post-infectious syndrome (e.g., Multisystem Inflammatory Syndrome in Children; MIS-C).
2. In surveillance and epidemiologic studies.

2.3. Clinical spectrum of COVID 19

WHO has classified COVID-19 disease in the following three categories:

1. **Critical COVID-19**– Defined by the criteria for acute respiratory distress syndrome (ARDS), sepsis, septic shock, or other conditions that would normally require the provision of life-sustaining therapies such as mechanical ventilation (invasive or non-invasive) or vasopressor therapy.
2. **Severe COVID-19**– (Moderate cases are also included in this category)
Defined by any of the following:
 - Oxygen saturation < 90% on room air*;

- Respiratory rate ≥ 60 breaths/min in children < 2 months old; ≥ 50 in children 2–12 months old; ≥ 40 in children 12–60 months old; and > 30 breaths/min in children > 60 months old;
- Signs of severe respiratory distress (accessory muscle use, inability to complete full sentences, and, in children, very severe chest wall in drawing, grunting, central cyanosis, or presence of any other general danger signs).

3. Non-severe COVID-19– Defined as absence of any criteria for severe or critical COVID-19.

*Saturation measurement in children is very difficult and unreliable, clinical judgment is very essential to treat a child with SpO_2 monitoring. So, in a clinical setting $\text{SpO}_2 < 94\%$ should be meticulously monitored and if needed aggressive treatment as severe COVID-19 should be done.

***Acute respiratory distress syndrome**

- Respiratory symptoms of hypoxemia and radiological change within 1 week of disease onset due to SARS-CoV-2 not explained by acute left ventricular failure or volume overload.
- The radiological findings of new infiltrates consistent with acute pulmonary parenchymal disease
- Partial pressure of oxygen (PaO_2) to fraction of inspired oxygen (FiO_2) ratio to be used as measure of oxygenation for noninvasive ventilation and a oxygenation index for invasive mechanical ventilation.
- Wherever partial pressure of oxygen (PaO_2) measurement facility is not available oxygen saturation to fraction of inspired oxygen (FiO_2) ratio to be used for the noninvasive ventilation and oxygenation saturation index (OSI) for invasive ventilation.

- The Severity of ARDS is given in the following table:

Oxygenation	Non-Invasive mechanical Ventilation	Invasive mechanical ventilation		
	No severity stratification	Mild	Moderate	Severe
	Face mask bi-level ventilation or AP CPAP ≥ 5 cm H ₂ O PF ≤ 300 SF ratio ≤ 264	$4 \leq \text{OI}^* < 8$ $5 \leq \text{OSI}^* < 7.5$	$8 \leq \text{OI} < 16$ $7.5 \leq \text{OSI} < 12.3$	$\text{OI} \geq 16$ $\text{OSI} \geq 12.3$

*OI = $(\text{FiO}_2 \times \text{mean airway pressure} \times 100) / \text{PaO}_2$

#OSI = $(\text{FiO}_2 \times \text{mean airway pressure} \times 100) / \text{SpO}_2$

*Rapid respiration (age based): < 2 months $\geq 60/\text{min}$;
2-12 months $\geq 50/\text{min}$;
12-60 months $\geq 40/\text{min}$;
>60 months $> 30/\text{min}$

2.4 Management of children with COVID-19 infection

2.4.1 Outpatient management of Children with COVID-19

The parents should be counseled regarding isolation, monitoring for clinical deterioration, and supportive care. They must be informed about the chances of clinical deterioration, which may occur suddenly after one week of symptoms.

The symptoms of clinical deterioration include:

- Severe respiratory distress,
- Difficulty breathing (for infants: grunting, central cyanosis, inability to breastfeed),

- Chest pain or pressure,
- Blue lips or face,
- Findings associated with shock (e.g. cold, clammy, mottled skin; new confusion; difficulty arousing; substantially reduced urine output), etc.

Use of telephone and digital media like video call can be used for the purpose of counseling and monitoring.

2.4.2 Duration of home isolation:

At 10 days after illness onset, recovery of replication-competent virus in viral culture is decreased and approaches zero.

For persons recovered from COVID-19 illness, the CDC recommends that isolation be maintained for **at least 10 days after illness onset and at least 3 days (72 hours) after recovery**. Illness onset is defined as the date symptoms begin. Recovery is defined as resolution of fever without the use of fever-reducing medications with progressive improvement or resolution of other symptoms.

A. Non-severe disease

- Children to remain in home isolation under parental supervision.
- These children should be given symptomatic treatment such as antipyretics for fever and pain, adequate nutrition and appropriate rehydration.
- Antibiotic therapy or prophylaxis should not be used in these children.
- COVID appropriate behavior including face masks (for more than 5 year of age, hand hygiene and physical distancing advised.
- **Any routine medications, which the child is taking, and vaccination should be continued.**
- Caregivers of children with COVID-19 should also monitor their patients for any signs and symptoms of clinical deterioration requiring an urgent re-evaluation. These include difficulty breathing/fast or shallow breathing, blue lips or face, chest pain or pressure, new confusion as well as an inability to wake up, interact when awake, drink or keep liquids down. For infants these include grunting and an inability to breastfeed.

- Home pulse oximetry measurements should be done regularly to look for any deterioration or desaturation so that they bring their child to the hospital if saturation drops below 94%.

2.5 Indication for hospital admission:

1. Children with severe or critical COVID-19
2. Children with non-severe COVID-19 may require hospital admission if they are at risk for severe disease due to underlying conditions (e.g. immunocompromised).

2.5.1 Indications for admission to the hospital for non-severe disease

- Presence of certain co-morbid conditions that lead to rapid clinical deterioration in a child with non-severe disease require hospital admission which include:
 - Diabetes
 - Congenital heart disease
 - Chronic lung disease
 - Cerebrovascular disease
 - Chronic kidney disease
 - Nephrotic syndrome
 - Immunosuppression
 - Obesity
 - Malignancy

Other patients can be managed in home under telemedicine guidance of the Health Care provider. Oral antimicrobials to be used on a case by case basis, especially for children with high risk of disease progression (immunocompromised, congenital heart disease, malignancy, etc.) and also in those cases with suspicion of bacterial co-infection (after sending blood culture) as per the hospital antibiotic policy. Oral Paracetamol and other symptomatic treatment to be done as explained above. Rest to be done as per treatment of non-severe infection.

2.5.2 Indication for referral to higher centers:

1. Children with COVID-19; worsening severe or critical disease
2. Children with non-severe COVID-19 who are at risk for severe disease due to underlying conditions (e.g. immunocompromised).

2.6 Investigations (as per clinical judgment and availability)

Although not all children infected with COVID-19 require investigations, they are mandatory for all children with severe and critical disease.

A. For those children with non-severe disease on case to case basis:

- Complete blood count,
- Serum electrolytes (sodium, potassium), blood glucose,
- Erythrocyte Sedimentation rate (ESR),
- Blood culture (if secondary infection is suspected),
- Chest X-ray

B. For Severe Cases (as per clinical judgment and availability of the tests)

- Complete Blood Count,
- Serum electrolytes (sodium, potassium),
- Blood glucose,
- Erythrocyte Sedimentation rate (ESR),
- C-Reactive Protein,
- Liver Function Test,
- Renal Function Test (serum urea, creatinine),
- Liver Function test,
- Inflammatory markers: Serum ferritin, D-dimer, serum Lactate dehydrogenase (LDH),
- Coagulation profile: aPTT, PT/INR,
- Arterial Blood Gas (ABG),

- Blood culture and sensitivity (to rule out bacterial sepsis/coinfection),
- Chest- X ray
- CRP, D-dimer, ferritin, and LDH needs to be done two or three times per week to monitor for cytokine storm

2.7 Management of COVID-19 with Severe and Critical infection:

To be admitted in COVID High Dependency Unit (HDU) or Pediatric Intensive Care Unit (PICU). The treatment strategies involve use of

- Supplemental oxygen,
- Fluid and electrolyte balance,
- Use of corticosteroids (as indicated),
- Low molecular heparin (as indicated),
- Management of ARDS,
- Sepsis, shock,
- Renal replacement Therapy (as per the standard guidelines) nutrition,
- Avoidance of unproven drugs.

2.7.1 Supplemental Oxygen

Indications for supplemental oxygen:

- Children with Oxygen Saturation less than 94% in room air,
- Obstructed or absent breathing,
- Severe respiratory distress,
- Central cyanosis,
- Shock,
- Coma or convulsions.

While interpreting the oxygen saturation level we should take into consideration about the child's previous disease condition and baseline SpO_2 e.g. a child with chronic lung disease or congenital heart disease might have a baseline of SpO_2 of 90%, which can be considered normal for that child. So, a cutoff of 94% can be used in previously healthy children without underlying disease conditions.

Modes of giving oxygen therapy:

Non-invasive	Invasive
Face mask	Intubation and Invasive Mechanical Ventilation
Nasal cannula	HFOV
Hood box (Head box)	Extracorporeal Membrane Oxygenation (ECMO)
Non-rebreathing mask	
HFNC (High-Flow Nasal Cannula)	
NIPPV (Noninvasive Positive Pressure Ventilation)	

A. Heated Humidified HFNC (High-Flow Nasal Cannula):

Indicated in children with COVID-19 who persist to have increased work of breathing and hypoxemia on supplemental oxygen

- Flow rate for HFNC therapy is same for all children regardless of disease conditions.
 - o < 12 kg: 2 L/kg/min
 - o >12 kg: 2 L/kg/min + 0.5 L/kg/min for each kg thereafter (max flow 50 L/min)
 - o Increase flow to the prescribed rate over a few minutes as tolerated.
 - o When supplemental oxygen is required, titrate FiO_2 to maintain the target SpO_2 .
- Airborne precautions to be followed with adequate PPE. HFNC should be tried for a maximum of 1-2 hours.
- Signs of improvement are decrease in heart rate and respiratory rate by 10-20%, decrease in FiO_2 requirement to less than 50% and improvement in oxygen saturations.
- Patients with progressive respiratory distress despite HFNC, or where HFNC is unavailable, can be escalated to NIV, bCPAP, or bi-level positive

airway pressure (BiPAP). Patients with worsening hypercapnia, acidemia, respiratory fatigue, hemodynamic instability or those with altered mental status should be considered for early invasive mechanical ventilation.

B. Non-Invasive Ventilation:

- Routine use of NIV is not recommended in COVID-19. It should be used only in selected patients with hypoxemic respiratory failure (**mild cases of ARDS without hemodynamic instability but needs close monitoring for escalation of treatment.**)
- Ideally, negative pressure single rooms are preferable for patients on NIV; in lack of such rooms keeping a distance of at least two meters between two beds should be considered.
- Conventional ventilators with NIV option having double lumen tubing is a safer option than NIV ventilator with single lumen tubing requiring exhalation port to washout the CO_2 .
- Antiviral/Antibacterial filters should be attached to the exhalation limb of the circuit to reduce environmental contamination. Preferred interfaces are helmet (hood), total face mask and oro-nasal non-vented masks.
- $\text{PaO}_2/\text{FiO}_2$ is a sensitive and accurate indicator of oxygenation function on NIV and can be used to define the severity of ARDS once the patient has been on a PEEP of 5 cm or more for a minimum of 30 minutes.
- Invasive ventilation must be considered if $\text{PaO}_2/\text{FiO}_2$ ratio is below 300.
- NIV might reduce intubation and mortality in mild ARDS, it is associated with higher mortality in moderate-to-severe ARDS.

C. Bubble CPAP:

- In resource limited settings, bubble CPAP should be considered for respiratory support in children with hypoxemia, severe pneumonia and/or ARDS where both non-invasive and invasive mechanical ventilation are not available.
- Bubble nasal CPAP (commercial or indigenous) may be used for newborns and children with severe hypoxemia as these are readily available alternatives in resource-limited settings.
- For minimization of environmental contamination, the infant could be placed in an oxygen hood to reduce droplets.
- These patients should be on continuous monitoring and in case the patient acutely deteriorates or does not improve after a short trial (about 2 hours) the patient needs to be intubated.

D. Endotracheal Intubation and mechanical ventilation:

Indications for invasive mechanical ventilation:

- Moderate/severe ARDS with $\text{PaO}_2/\text{FiO}_2$ ratio below 200,
- Hemodynamic instability
- Multi-organ failure
- Abnormal mental status
- Patients with worsening hypoxia and work of breathing

2.7.2 Corticosteroids:**Indications for using glucocorticoids:**

1. Children with severe or critical COVID-19 who require mechanical ventilation or those who require supplemental oxygen and have risk factors for disease progression or those with rapid disease progression or those who develop septic shock.

2. Steroids should be continued in children with an underlying condition requiring chronic steroid treatment.

3. Cases of comorbid conditions where steroid therapy is indicated like Nephrotic Syndrome (NS). Steroids are to be used only after, first 3-5 days of illness, as it may prolong viral clearance, if used earlier. Low-dose glucocorticoid regimens include one of:

- **Dexamethasone** 0.15 mg/kg orally, IV, or through nasogastric tube (NG tube) once daily (maximum dose 6 mg) OR
 - **Prednisolone** 1mg/kg orally or NG once daily (maximum dose 40mg) OR
 - **Methylprednisolone** 0.8 mg/kg IV once daily (maximum dose 32 mg)
- OR

- **Hydrocortisone**

- o For neonates (<1 month of age): 0.5 mg/kg IV every 12 hours for 7 days followed by 0.5 mg/kg IV once daily for 3 days;
- o For children ≥ 1 month: 1.3 mg/kg IV every 8 hours (maximum dose 50 mg; maximum total daily dose 150 mg).

The duration of therapy is 5-7 days.

2.7.3. Low Molecular Weight Heparin:

LMW heparin is known to reduce the risk of Venous Thromboembolism (VTE) and may have anti-inflammatory properties. Unlike adults, the decision to start venous thromboembolism (VTE) prophylaxis in children is individualized.

A. Indications for starting prophylactic Low Molecular Weight Heparin in children:

1. Strong personal or family history of VTE
2. Indwelling central venous line and two or more additional risk factors
3. Four or more risk factors

Risk factors for thrombosis to consider:

- Personal history of thrombophilia or VTE
- First degree relative with VTE
- Presence of central venous line
- Congenital Heart Disease
- Post pubertal age
- Prematurity
- Antiphospholipid syndrome
- Decreased mobility from baseline
- Burns
- Active malignancy
- Indications of venous stasis or cardiac low flow state
- Estrogen therapy
- Active systemic infection
- Flare of inflammatory disease
- Obesity
- Severe dehydration
- Recent surgery or trauma

An assessment of bleeding risks (intracranial hemorrhage, active bleeding, coagulopathy, neurosurgical procedure within 24 hour, etc.) versus benefit should be compared on each pediatric patient. Alternative methods of prophylaxis, such as early ambulation or mechanical prophylaxis should be considered in contraindicated patients and all COVID-19 pediatric patients, if applicable.

B. Dose and route of prophylactic Enoxaparin:

0.5 mg/kg (maximum dose 40 mg) given twice a day subcutaneously.

It can be administered alternatively between the left and right anterolateral and left and right posterolateral abdominal wall. To avoid bruising, the injection site should not be rubbed.

Monitoring of patients on Enoxaparin: If facilities are available in the center, anti-factor Xa levels after 4 hour of subcutaneous dose completing the third consecutive dose can be done to obtain a level of 0.20-0.49 anti-Xa U/mL.

C. Indication for therapeutic anticoagulation:

- Documented or strongly suspected venous thromboembolism (VTE)
- Clotting of vascular access devices.
- Patients receiving anticoagulation therapy prior to admission.

For such children injection enoxaparin is given at the dose of **2 mg/kg subcutaneously** every 12 hourly with an anti-Xa factor target of 0.5–1 IU/ml.

Individuals who have suffered from venous thromboembolism require three months of anticoagulation after they get discharged.

2.8 Management of COVID-19 with ARDS (CARDS)

A. Mild ARDS: High flow nasal oxygen or non-invasive ventilation (BiPAP or CPAP) may be given with close monitoring for increase in severity of illness.

B. Moderate to Severe ARDS: The following lung protective ventilation strategy to follow:

Parameter	Strategy
Low Tidal Volume	4-8 ml/kg of predicted body weight
Plateau Pressure	< 28 cm H ₂ O
Mean Airway Pressure	<18-20 cm H ₂ O
Driving pressure	<15 cm H ₂ O
Positive End Expiratory Pressure (PEEP)	Moderate ARDS < 10 cm H ₂ O Severe ARDS 10-15 cm H ₂ O
FiO ₂	<60%
Sedation with or without neuromuscular blockade	Adequate
ET tube	Cuffed
Suctioning	Inline suctioning preferred to open suctioning (to avoid disconnections)
Nutrition	Enteral nutrition within 24 hour
Position	Prone (ideally 12-16 hours per day, only if feasible in severe ARDS)
Fluids	Restrictive fluid strategy, 2/3rd of total maintenance fluid
Weaning	Daily assessment for weaning, early weaning.

Blood Transfusion	If oxygenation and hemodynamically stable, threshold Hb <7; if hypoxemia or shock, threshold Hb<10.
-------------------	---

High-frequency Oscillatory Ventilation (HFOV) (for refractory respiratory failure), Extracorporeal Membrane Oxygenation (ECMO) may be considered in patients with continued severe hypoxemia despite maximal support if facilities is available.

2.9 Management of shock

- Give Intravenous fluid bolus **10–20 ml/kg per bolus up to 40–60 ml/kg**, over the first hour of resuscitation.
- The initial fluid of choice should be **crystalloids**. In children with COVID-19 and shock, age-appropriate mean arterial pressure (MAP) should be targeted. In settings where accurate MAPs cannot be easily obtained, systolic blood pressure is an acceptable option. For inotropic support, **epinephrine** should be chosen as the first-line vasoactive infusion. **Epinephrine should be considered as the first-line agent in patients with shock and if not improved other inotropes can be added. For further management NEPAS guidelines for Shock can be referred to.**
- If there is no improvement with one inotrope another one can be added accordingly. Vasopressin should be considered in children who need high doses of catecholamines. Inodilators such as milrinone, dobutamine or levosimendan could be used when there are signs of tissue hypoperfusion and cardiac dysfunction, despite high doses of catecholamines.
- Hydrocortisone should be added if there is fluid refractory catecholamine resistant shock (avoid if already on dexamethasone or methylprednisolone).
- Once stabilized, restrict IV fluids to avoid fluid overload. Enteral nutrition to be initiated as early as possible if there is no contraindication. Empirical antibiotics to be instituted within the first hour after sending the blood culture, as per the hospital policy.

2.10 Empiric antibiotics:

For patients with suspected or confirmed asymptomatic COVID-19, antibiotic therapy or prophylaxis is not warranted. For those patients with non-severe COVID-19, antibiotics should not be prescribed unless there is clinical suspicion of a bacterial infection. For patients with suspected or confirmed severe COVID-19, the use of empiric antimicrobials to treat all likely pathogens is recommended, based on clinical judgment, patient host factors and local epidemiology, and this should be done as soon as possible (within 1 hour of initial assessment if possible), ideally with blood cultures obtained first. Antimicrobial therapy should be assessed daily for de-escalation. Duration of empiric antibiotic treatment should be as short as possible; generally 5–7 days.

2.11 Antiviral therapy, Remdesivir:

WHO does not recommend the use of Remdesivir in children. It may be used in case-to-case basis as per clinical judgment of the treating pediatrician.

Remdesivir is dosed according to weight as follows:

- ≥ 3.5 to < 40 kg: 5 mg/kg intravenous (IV) loading dose on day 1, followed by 2.5 mg/kg IV every 24 hours
- ≥ 40 kg: 200 mg IV loading dose on day 1, followed by 100 mg IV every 24 hours

The usual duration of therapy is up to 5 days for children with severe disease; for children with critical disease who are not improving after 5 days, the duration may be extended to up to 10 days.

2.12 Renal Replacement therapy:

Whenever it is indicated as per protocol.

2.13 When to discontinue isolation in admitted children?

- For asymptomatic persons: 10 days after testing positive.
- For severe, symptomatic and immunocompromised children, the isolation days may be extended as per infectious disease experts opinion.

CHAPTER III

NEONATES WITH COVID-19

The immature immune system, passive transfer of maternal IgG antibodies, and lower ACE-2 expression may result in less inflammation, milder illness, and hastened recovery in infants and children compared to adults.

3.1 Early-onset neonatal COVID-19:

This manifests between 2 and 7 days after birth. Most neonates are asymptomatic or have:

- Mild symptoms such as rhinorrhea and cough and fever
- Moderate to severe symptoms such as respiratory distress, poor feeding, lethargy, vomiting and diarrhea and rarely clinical evidence of multi-organ failure have been observed.

Laboratory finding

There might be leukocytosis, lymphopenia, thrombocytopenia, and non-specific findings of elevated inflammatory markers.

Management:

The management is supportive.

3.2 Late-onset neonatal COVID-19:

This manifests beyond 5 to 7 days after birth. This might have been acquired from respiratory secretions of mother, caregivers, or household contacts. Many affected neonates have negative initial RT-PCR test results (at 24 and 48 hours after birth) before initial discharge from the hospital and get re-admitted with symptoms suggestive of COVID-19.

Clinical Features:

- Hyperthermia
- Coryza
- Mild respiratory symptoms
- Apnea
- Poor feeding or vomiting
- Lethargy

Age less than 1 month has been associated with a 3-fold higher risk of critical care admission.

Management:

Management includes supplemental oxygen, respiratory support, fluid resuscitation, and temperature control. Antiviral medications and steroids in neonatal COVID-19 are lacking. However use of Remdesivir, dexamethasone has been reported.

3.3 Multisystem inflammatory syndrome in neonates (MIS-N):

Neonatal MIS (MIS-N) has rarely been reported. Neonates are usually asymptomatic or they have mild illness but some of them may develop severe symptoms. This entity in neonates may be under reported as these babies might present with signs of sepsis. Hence, MIS-N diagnosis should be considered on babies presenting with multisystem involvement and evaluated accordingly.

3.4 Prevention of transmission from infected mother to newborns

3.4.1 Key points in neonatal resuscitation

The following points must be noted:

- Use of appropriate PPE by the caregivers.

- Initial steps are unlikely to be aerosol generating; they include drying, tactile stimulation, placement into a plastic bag or wrap, assessment of heart rate, and placement of pulse oximetry and ECG leads.
- Suction should not be performed unless indicated.
- Endotracheal medications should be avoided.
- Intravenous epinephrine via a low-lying umbilical venous catheter is preferred.
- Closed incubators should be used.

3.4.2. Testing in neonates

- Testing for SARS-CoV-2 RNA by RT-PCR should be done for all neonates born to COVID positive mothers even in the absence of signs of infection in the neonate.
- Obtain either a single swab of the nasopharynx or a single swab of the throat followed by the nasopharynx or two separate swabs from each of these sites for RT-PCR test.
- Testing for both symptomatic and asymptomatic neonates born to suspected or confirmed COVID-19 should be done first at approximately 24 hours of age and if initial test results are negative, or not available, testing should be repeated at 48 hours of age.
- If it is planned that a healthy newborn will be discharged prior to 48 hours of age, clinicians may choose to order a single test at 24-48 hours of age.
- For infants who require ongoing hospital care, caregivers should continue to use appropriate PPE until discharge, or until the infant has two consecutive negative tests collected ≥ 24 hours apart.

3.4.3. Mother-baby contact

- Babies born to suspected or confirmed COVID-19 should be enabled to remain together and practice skin-to-skin contact.
- Breastfeeding should be established early and mothers should be encouraged to continue breastfeeding.

- The newborn's risk for acquiring SARS-CoV-2 from the mother is low, and data suggest no difference in risk of neonatal SARS-CoV-2 infection whether the neonate is cared for in a separate room or remains in the mother's room.
- Rooming-in helps establish breastfeeding, facilitates bonding and parental education, and promotes family-centered care.
- As there is a potential risk of SARS-CoV-2 transmission to the neonate via contact with infectious respiratory secretions from mother or caregivers, all caregivers should practice infection prevention and control measures while caring for a neonate.
- If separation is necessary for mothers who are too sick to care for their sick babies or who need high levels of care, expressed breast milk should be given to babies.
- If a baby's RT-PCR is positive for COVID-19, separation of mother with suspected or confirmed COVID from baby in order to reduce the risk of transmission of disease is not required.
- Measures to reduce the risk of transmission:
 - o Mothers or caregivers should wear a mask and practice hand hygiene during contact with neonates.
 - o Masks and face shields are not recommended for neonates or children < 2 years of age.
 - o Maintain a physical distance of >2 meter between mother and neonate in other times if feasible.

3.5. Discontinuation of isolation and precaution guidelines for confirmed COVID mothers

- At least 10 days have passed since symptoms first appeared (up to 20 days if they have more severe to critical illness or are severely immuno compromised).
- At least 72 hours have passed since their last fever without the use of antipyretics.
- Other symptoms have improved.
- For asymptomatic mothers, after at least 10 days of positive test results.

3.6. Breastfeeding:

- The risk of SARS-CoV-2 transmission from ingestion of breast milk is minimal.
- Mothers with suspected or confirmed COVID-19, should be encouraged to continue breastfeeding and she should be counselled for the benefits of breastfeeding, which outweigh the potential risks of transmission.
- A breastfeeding mother who is not fully vaccinated against COVID-19 should take precautions to protect themselves and the breastfed child when either member of the family has suspected or confirmed COVID-19.
- Mothers and caregivers should follow these precautions during their recommended period of isolation:
 - o Wash hands with soap and water before touching the baby or expressing breast milk.
 - o Use of hand sanitizer if soap and water are not available.
 - o Wear mask
- If a mother is not able to breastfeed her baby due to her severe illness, she should be encouraged to express milk for her baby following appropriate IPC measures.
- If a mother is very sick to breastfeed or express breast milk, we need to discuss with mother and parents regarding donor human milk based on the local culture and practice.
- If this is also not possible, consider wet nursing (defined as another woman breastfeeds the child) or appropriate breast milk substitutes, informed by feasibility, safety, sustainability, cultural context, acceptability to mother and service availability.

3.7. SARS-CoV-2 vaccines safety in lactating mothers

Lactating mothers should receive a vaccine against SARS-CoV-2. COVID-19 antibodies present in breast milk may have protective effects in infants.

Refer to Flowchart 1, 2, and 3 for an approach to neonates born to suspected and positive mothers in Appendices.

CHAPTER IV

MULTI-SYSTEM INFLAMMATORY SYNDROME IN CHILDREN (MIS-C)

4.1. Background

Multisystem Inflammatory Syndrome in Children (MIS-C) also known as pediatric hyper-inflammatory syndrome, pediatric multisystem inflammatory syndrome (PIMS) or pediatric hyper-inflammatory shock. MIS-C is a condition that is temporally associated with SAR-CoV-2 and is characterized by fever, inflammation and multi-organ dysfunction, which manifests several weeks after the infection of SARS-CoV-2. This condition should be differentiated from other pediatric inflammatory conditions like Kawasaki disease, bacterial sepsis, staphylococcal and streptococcal toxic shock syndromes and macrophage activation syndromes as these conditions have features similar to that of MIS-C.

4.2. When to suspect MIS-C?

Any children who fulfil the case definition criteria given by CDC or WHO should be evaluated for MIS-C.

WHO case Definition of MIS-C

- Children and adolescents (0-19 years) with fever \geq 3days

AND: Clinical sign of multisystem involvement (at least 2 of the following):

1. Rash or bilateral non-purulent conjunctivitis or muco-cutaneous inflammation signs (oral, hands or feet).
2. Hypotension or shock
3. Features of myocardial dysfunction, pericarditis, valvulitis, or coronary abnormalities (including ECHO findings or elevated Troponin/NT-pro BNP)

<p>4. Evidence of coagulopathy (by PT, PTT, elevated D-dimers)</p> <p>5. Acute gastrointestinal problems (diarrhea, vomiting, or abdominal pain)</p>
AND
<ul style="list-style-type: none"> Elevated ESR, C-reactive protein, or procalcitonin
AND
<ul style="list-style-type: none"> No other obvious microbial cause of inflammation, including bacterial sepsis, staphylococcal or streptococcal shock syndromes
AND
<ul style="list-style-type: none"> Evidence of COVID-19 (RT-PCR, antigen test or serology positive), or likely contact with patients with COVID-19

Diagnostic evaluation of MIS-C

<p>A child suspected for MIS-C should also be evaluated for other common clinical disease conditions like infectious and non-infectious causes that justify the patient's clinical manifestation.</p>
<p>Outpatient evaluation for MIS-C may be appropriate for assessing well-appearing children with stable vital signs and for ensuring that physical examinations provide close clinical follow-up.</p>
<p>Patients presenting with shock, significant respiratory distress, neurologic changes (altered mental status, encephalopathy, focal neurologic deficits, meningismus, and papilledema), dehydration, or features of Kawasaki Disease (KD) should be admitted for further evaluation, regardless of MIS-C status, in accordance with standard of care.</p>
<p>Refer to Appendix 3 I for clinical manifestations and details of investigations that can be done for MIS-C.</p>
<p>Refer to Appendix 3 II for distinctions between MIS-C and Kawasaki disease.</p>
<p>Refer to Appendix 3 III for definition of organ system involvement.</p>
<p>Refer to Appendix 2 - flow chart 4 and 5 for the evaluation and management of MIS-C.</p>

4.3. Indications for hospitalization

- Features of Kawasaki disease (KD)
- Severe abdominal pain or vomiting, especially if unable to tolerate oral feeds
- Clinical or laboratory evidence of dehydration
- Marked elevations of inflammatory markers (CRP >10 mg/dl)

4.4. Indications for PICU/HDU admission (if facilities available)

- Abnormal vital signs (tachycardia, tachypnea)
- Respiratory distress of any severity
- Abnormal EKG findings
- Hemodynamic instability (shock, arrhythmia)
- Significant respiratory compromise/ Severe respiratory distress/ respiratory failure
- Evidence of cardiac involvement (myocarditis, elevated troponin or brain natriuretic peptide, depressed ventricular function or coronary artery abnormality on echocardiogram, abnormal echocardiogram)
- Neurologic changes (e.g., depressed mental status, abnormal neurologic examination, seizures)
- Evidence of acute kidney injury, acute hepatic injury, or coagulopathy
- Underlying medical condition that may place the child at increased risk for complications (e.g., immunodeficiency, cardiac or pulmonary conditions)

4.5. Approach to a child with suspected MIS-C

Refer to Flow chart 4 in the appendices

4.6. Management of MIS-C

Refer to Flow chart 5 in the appendices

4.7. Treatment protocols can be grouped under 4 major categories

1. Supportive care
2. Antibiotics
3. Cardiac management
4. Immunomodulatory treatment
5. Antiplatelet and anticoagulation therapy

1. Supportive care

- Primary Assessment Pentagon (ABCDE).
- The ABCDE approach consists of stabilization of airway, breathing, circulation and neurological status.
- If the patient is in shock, vasoplegic or cardiogenic, manage with fluids, Vasopressors or inotropes as per the protocol.

2. Antibiotics

- Empirical first dose of broad-spectrum antibiotics of ceftriaxone and vancomycin/ cloxacillin preferably within the first hour of presentation after blood culture is obtained.
- Clindamycin is added if there are features consistent with toxin-mediated illness (e.g., erythroderma). Antibiotics should be discontinued once bacterial infection has been excluded if the child's clinical status has stabilized.

3. Cardiac management

- BNP and/or troponin T at diagnosis should be trended over time until they normalize.
- EKGs every 48 hours and during follow up visits for detecting conduction abnormalities.
- Echocardiograms at diagnosis and follow up for ventricular/valvular function, pericardial effusion, coronary artery (CA) dimensions using z-scores.

- Echocardiograms repeated at a minimum of 7-14 days and 4-6 weeks after initial presentation. Echo should be repeated 1 year after MIS-C diagnosis in children with cardiac involvement during acute phase of illness. Children with LV dysfunction and coronary artery aneurysms (CAAs) will require more frequent echocardiography.
- Cardiac MRI at 2-6 months post-acute illness may be considered in children with moderate to severe LV dysfunction to evaluate for myocardial fibrosis and scarring.

4. Immunomodulatory treatment

- Patients with mild symptoms and without organ dysfunction may only require close monitoring and supportive treatment without immunomodulatory treatment.
- Stepwise progression of immunomodulatory therapies to be used with IVIG as 1st line of therapy.
- IVIG should be given to MIS-C patients who are hospitalized and or fulfill KD criteria.
- High-dose IVIG (typically 2 gm/kg, based on ideal body weight) should be used for treatment of MIS-C.
- Before starting IVIG, cardiac function and fluid status should be assessed. Children with depressed cardiac function will require close monitoring and diuretics while IVIG administration.
- In some patients with cardiac dysfunction, IVIG may be given in divided doses (1 gm/kg daily over 2 days).
- IV Methylprednisolone (MP) 1–2mg/kg/day should be given with IVIG as adjunctive therapy for patients with shock and/or organ dysfunction.
- In patients who do not respond to IVIG + MP and requires high dose or multiple inotropes and/or vasopressors, high-dose, IV pulse Methylprednisolone (10–30 mg/kg/day) may be considered.
- In patients with refractory MIS-C despite a single dose of IVIG, a second dose of IVIG is not recommended, given the risk of volume overload and hemolytic anemia associated with large doses of IVIG.
- IV MP (1–2 mg/kg/day) may also be considered with milder forms of MIS-C who are persistently febrile and symptomatic despite a single dose of IVIG.

- Anakinra (>4 mg/kg/day IV or SC) may be considered in patients with features of macrophage activation syndrome, when long-term use of glucocorticoids is contraindicated, or treatment of MIS-C is refractory to IVIG and glucocorticoids.
- If IVIG is not available, MP (2mg/kg/day in 2 divided doses) should be given.
- After defervescence and clinical improvement, steroids can be changed to an equivalent oral dose of Prednisolone at discharge and tapered off over 3-4 weeks.
- Other immunomodulation: Anakinra, Tocilizumab, Infliximab.

5. Antiplatelet and anticoagulation therapy

5.1. When to seek help?

There are 3 ways in which stress can affect everybody including the C&A. Sometimes these problems can make doing day-to-day life very difficult. This could be because of:

- Problems with feelings (Emotional symptoms): e.g. feeling sad, angry, irritated, crying, fearful and anxious.
- Problems in the body (Physical symptoms): e.g. headaches, abdominal pain, body aches, fainting like episodes, trembling, weakness of limbs, etc.

Low dose aspirin (3-5mg/kg/day; max 81mg/day)

Indications

- All cases with diagnosis of MIS-C
- MIS-C with CAAs and a maximal z-score of 2.5-10.0

Contraindications

- Active bleeding
- Significant bleeding risk
- Platelet count $\leq 80,000/\mu\text{L}$

Duration: until normalization of platelet count and confirmed normal coronary arteries at ≥ 4 weeks after diagnosis

Enoxaparin: subcutaneous dose

- < 2months: 1.5 mg/kg/dose every 12 hours (therapeutic), 0.75 mg/kg/dose every 12 hours (prophylactic)
- >2months: 2 mg/kg/dose every 12 hours (therapeutic), 0.5 mg/kg/dose every 12 hours (prophylactic)
- MIS-C with CAA z score ≥ 10.0 should be treated with low dose aspirin and enoxaparin (factor Xa level 0.5-1.0) or warfarin

- Longer outpatient enoxaparin dosing
- CAAs with z-score of >10.0 (indefinite treatment)
- Documented thrombosis (treatment for >3 months pending thrombus resolution)
- Ongoing moderate to severe LV dysfunction

CHAPTER V

CRISIS RESPONSES AND PSYCHO-SOCIAL MANAGEMENT FOR CHILDREN AND ADOLESCENTS (C&A) DURING COVID-19 PANDEMIC

- Problems in our behavior/action (Behavioral symptoms): e.g. Irritation, clinginess to parents, excessive crying, aggression, demanding behaviors, substance use, school refusal, etc.
- There can be problems in sleep and appetite (Eating habits).
- The caregivers should seek help if the C&A present with the above-mentioned symptoms that are well in excess than normal and haven't responded to normal intervention. Symptoms are severe enough to cause disruption of daily functioning (even after applying coping techniques as mentioned in figure 6, then the child should be referred for further evaluation to a mental health professional.
- Pharmacotherapy for symptoms secondary to stress resulting in impairment in functioning and not improving after coping techniques.

I. For children below 6 years of age

- Pharmacotherapy is not the first choice
- If needed refer to a specialist (C&A Psychiatrist or Psychiatrist)

II. For children 6 to 12 years of age:

- Tablet SERTRALINE 25 mg ½ tablets PO in the morning after food.
- Can be increased tablet SERTRALINE 25 mg 1 tablet PO in the morning after food if needed.

III. For children more than 12 years of age:

- Tablet SERTRALINE 25 mg PO in the morning after food.
- Tablet CLONAZEPAM 0.25 mg PO HS (not more than 15 days)
- To follow up weekly for first 2 weeks then
- Follow up once fortnightly then
- Follow up monthly

•Note Medication not to be used for more than 6 months. If need arises to continue medication for more than 6 months, then refer to Child & Adolescent clinic or Psychiatrist.**

•For Assistance contact: 9808522410- CAP KCH Helpline and 16600110666 CAP KCH Hotline.

CHAPTER VI

MANAGING CHILDREN WITH DISABILITIES IN COVID-19 FACILITIES: GUIDANCE FOR DOCTORS, NURSES, PARAMEDIC AND OTHER HEALTH CARE STAFF

Children with disabilities need an innovative, different strategy to communicate, calm and engage them while they are admitted in ICU, isolation and quarantine facilities. It is imperative that health care workers need to adopt innovative and bespoke communication strategies to be able to effectively communicate with children with disabilities. The guidance below gives a few considerations for the health care staff which will help them to know and recognize the special needs of children with disabilities and enable them and the parent/caregiver to know that their differences are valued and supported. Children with disabilities may require bespoke support in communication, calming and creative engagement during their stay.

This guidance is about the key points to be considered when assessing and treating a child with learning disabilities or autism who is suspected of having or is known to have COVID-19. Other issues related to access to facilities, information, services, assistive devices, disability-specific therapy and therapeutics are general for all persons with disabilities and are not addressed in this guidance.

GENERAL:

1. Be aware of diagnostic overshadowing: It is important for the health care staff to differentiate the symptoms of physical ill health of children with disabilities and not to mistakenly attribute to a mental health or behavioral problem.

2. Information about the child's health needs, preferred method of communication and other preferences should be documented in the patient chart at admission by asking the child or the parent/caregiver by the nurse/paramedic.

3. Every child deserves and has the right to attain the highest standard of health care, without discrimination. It is also important not to make generalised judgements or assumptions about the vulnerability or frailty based on their dependence on others for support in daily living.

4. Involve and listen to the parent/caregiver as they have a wealth of information about the child with disabilities and how they have been, including any other comorbidities and the medication the person is taking. This needs to be documented in the child's chart.

5. You may seek support from rehabilitation experts (Speech & Language Therapist, Clinical Psychologists, Physiotherapists etc.) regarding communication, reasonable adjustments, and assessment of pain.

6. Presence of at least one regular parent/caregiver whilst the child with disabilities is in quarantine/isolation/hospital may be considered. This can be done if the parent/caregiver adheres to the required PPE and the institution policy.

COMMUNICATION:

1. Children with disabilities have the same illnesses as everyone else, but the way in which they respond to, or communicate their symptoms may be different and not obvious.

2. Check with the child and/or the parent/caregiver for the best way to communicate.

3. Use simple, clear language, avoiding medical terms and 'jargon' wherever possible. Some children with disabilities may be non-verbal and therefore unable to tell you how they feel. Communication boards, pictures, sign language may be a useful way of communicating.

4. A child with a learning disability, autism may not be able to articulate their response to pain in the expected way: for example,

a. they may say that they have a pain in their stomach when the pain is not there;

b. may say the pain is less acute than you would anticipate; or

c. not report experiencing pain when they are.

d. Some may feel pain in a different way or respond to it differently: for example, by displaying challenging behavior; laughing or crying; trying to hurt themselves; or equally may become withdrawn or quiet.

e. Children who use a wheelchair may have chronic pain.

5. Understanding what is 'normal' for the child is crucial to obtaining a full assessment and may aid the doctor reach a more accurate diagnosis.

CALMING & CREATIVE ENGAGING:

A change in the 'normal' daily routine or change in regular care giver may have a negative effect on the sick child with disabilities. This may happen in a hospital setting as well when there is a change in health care staff. Moreover, masks and protective clothing may frighten and, make them more anxious or lead to adverse behaviours, such as hurting other people, hurting themselves or damaging property. Do not assume that this is an indication of mental illness but do your best to work with the child, the parent/caregiver to find out how best to keep them calm and engaged.

The following toy kits will be of use in each COVID-19 facilities to calm the children who are distressed, anxious and agitated. And also, to creatively engage them during their institutional (hospital/isolation/quarantine) stay. They are simple, colourful toys, notebooks, crayon etc. which is commonly available in the market and/or the institution may consider allowing one toy/comforting piece of clothing that the child may be close to. (Refer to Fig: 9 and 10.)

CALMING TOY KIT

Purpose for use: Distraction play for calming the child's emotional distress

1. Squeeze Ball – squeeze and soothe stress/anxiety away (Tactile)
2. Light up Bouncy ball – Bounce and gaze at glitter (Visual, Tactile)
3. Rainbow Slinky Spring – Pull, drop down (Multisensory interactive)
4. Colourful Rattle – Shake it, roll it and hear it (Auditory)
5. Colourful pinwheel – Blow and see it spin (Multisensory Interactive)

ENGAGING TOY KIT

Purpose for use: Developmental play for stimulating brain functions

1. Notebook & Crayons – fine motor skills, creative expressions, conceptual learning
2. Building Blocks – eye-hand coordination, learning preschool concepts, visuconstruction play
3. Farm animals – make belief play, building language skills, concepts
4. Push bicycle toy – cause and effect , imitative & make belief play
5. Stickers – can be used as rewards

CHAPTER VII

INTEGRATING GENDER, EQUITY AND HUMAN RIGHTS (GER) IN CLINICAL MANAGEMENT (CM) OF COVID-19

Gender, equity and human rights matter in health and clinical management. The men and women, girls and boys, or any individual of gender and sexual identity experience differences in health status, exposure to risk and vulnerability, access to and use of services, health-seeking behaviour, experiences in health care settings, and health and social outcomes due to their biological and social standing in the society. Health inequities manifest in differential exposure, vulnerability, access, health outcomes and consequences, so it is very important to recognise these aspects and provide health services from gender, equity, and human rights perspectives. In response to COVID-19, the following aspects are suggested to strongly consider while managing COVID-19 patients in health facility settings.

1. PROVIDING RESPECTFUL CARE TOWARDS ALL PATIENTS

Naturally, we envision a relationship between patients and service providers characterized by caring, empathy, support, trust, confidence, and empowerment, as well as gentle, respectful, and effective communication to enable informed decision making. While dealing with the patients of COVID-19 and non-COVID-19 at clinical sites, health workers/providers need to be aware about the differences, providing fair treatment and respecting and protecting the rights of an individual. Health workers/providers need to:

- Demonstrate equal and fair treatment/ behavior irrespective of an individual's age, sex, caste, ethnicity, socioeconomic status, education, sexual orientation, family/ cultural background, disabilities or any other characteristics.
- Respect the right to information, informed consent and refusal; right to confidentiality, privacy, dignity, choices/preferences, equitable care; and self-determination; right to freedom from harm, ill treatment and discrimination; and right to timely healthcare and to the highest attainable level of health.

Managing social stigma:

Dos and Don'ts

DO – Use respectful and dignified verbal, and body language

Don't – Use offensive verbal and body language.*

DO - talk about the new coronavirus disease (COVID-19)

Don't - attach locations or ethnicity to the disease, e.g., “Chinese Virus”.

DO - talk about “people who may have COVID-19” or “people who are presumptive for COVID-19”

Don't - talk about “COVID-19 suspects” or “suspected cases”.

DO - talk about people “acquiring” or “contracting” COVID-19

Don't talk about people “transmitting COVID-19” “infecting others” or “spreading the virus” as it implies intentional transmission and assigns blame.

DO - speak accurately about the risk from COVID-19, based on scientific data and latest official health advice.

Don't - repeat or share unconfirmed rumours.

DO - talk positively and emphasise the effectiveness of prevention and treatment measures.

Don't - emphasise or dwell on the negative, or messages of threat.

DO - emphasise the effectiveness of adopting protective measures to prevent acquiring the coronavirus, as well as early screening, testing and treatment.

For more information:

[*National Guidelines for Disability Inclusive Health Services, 2019](#)
[covid19-stigma-guide.pdf](#)
[\(who.int\)](#)

- Avoid unintended biases towards women, girls, (with or without disabilities) or any clients based on their identity and socioeconomic backgrounds.
- Be aware of social stigma and Dos and Don'ts. Refer the box - there are some dos and don'ts on language¹ when talking about the COVID-19².
- Be aware of gender related biases, be non-judgmental.
- Be culturally sensitive and appropriate to age.
- Be aware of rights of childbearing women and respect those while providing care to them³.
- Be aware on data for men and women on predisposing factors, delays in seeking care, co-morbidities for the COVID-19 and risk groups and link to diseases as well as possible biological differences in COVID-19 impact over men and women. Even though COVID-19 infections are distributed equally among men and women, evidence shows more deaths among men due to biological factors, resulting presumptively due to a more robust immune response among women.
- Be aware in regards with non-health effects of COVID-19 pandemic. Evidence suggests women have borne the brunt of non-health impacts, including job and wage losses, increases in unpaid work in homes including health work, increases in violence against women, especially intimate partner violence, and lack of adequate social protection. They have also faced lack of access to needed non-COVID-19 health services, especially sexual and reproductive health services.

¹ Disability inclusive guidelines in English FINAL
 2covid19-stigma-guide.pdf (who.int)

³ Microsoft Word - Final_Respectful_Care_Charter_12-15-11.docx (who.int)

2. RESPONDING TO GENDER-BASED VIOLENCE (GBV)/ VIOLENCE AGAINST WOMEN AND GIRLS (VAWG)

The high prevalence of GBV/VAWG in Nepal is an ongoing challenge. NDHS 2016 reports 22% women experience physical violence. Global data show 1 in 3 women has experienced lifetime physical and/or sexual violence, mainly by an intimate partner. More importantly, available evidence points to significant increases GBV/VAWG increase in any emergency situation, and it has been exacerbated in COVID-19 situation and this has alarmed all actors working against GBV/VAWG.

What is gender - based violence (GBV)?

Gender based violence refers to harmful acts directed at an individual based on their gender. It is rooted in gender inequality, the abuse of power and harmful norms. GBV is a serious violation of human rights and a life-threatening health and protection issue. GBV is committed in many forms such as physical, emotional/psychological, sexual, cultural/social, economic or any kind that endangers the safety, health and well-being of an individual.

Domestic Violence refers to violent or aggressive behavior within home involving intimate partner and immediate family members.

The risks of violence that women and their children face during the current COVID-19 crisis cannot be ignored. “There never are excuses for violence”. Health systems have an important role in ensuring that services for survivors of gender-based violence remain accessible during the COVID-19 pandemic. The routine screening of GBV is NOT recommended by WHO during COVID-19 response. WHO guidance includes the dos and don'ts to be followed in this regard⁴. In case of a patient/client who comes to the health facility and discloses experience of violence, it is most important to respond⁵.

4 See 2014, WHO, UNW, UNFPA. Health care for women subjected to intimate partner violence or sexual violence: A clinical handbook: <https://www.who.int/publications/i/item/WHO-RHR-14.26>

5 See 2020 WHO: COVID-19 and violence against women: What the health sector / system can do: <https://www.who.int/reproductivehealth/publications/vaw-covid-19/en/>

Five Actions for Health workers/providers to respond to GBV/VAWG

1. Be aware of the increased risk and health consequences of GBV/VAWG in the context of COVID-19.
2. Recognize the signs and know when and how to ask about violence.
3. If violence is disclosed, act to provide timely care for physical, sexual, reproductive and mental health.
4. If violence is disclosed, provide First-line support and medical care to survivors. The first-line support is most important, and it involves 5 simple tasks of LIVES:
 - LISTEN: listen to women, girls closely, with empathy, and without judging.
 - INQUIRE: assess, identify and respond to person's various needs and concerns.
 - VALIDATE: show that you understand survivor's experience, feeling and believe her.
 - ENHANCE SAFETY: discuss a plan to protect the survivor from further harm if violence occurs again.
 - SUPPORT: support her by helping her connect to information, services, and social support.
5. Share information about available support, identify referral path ways and refer to other essential services.

Health facilities can identify and provide information about services available locally (e.g. hotlines, shelters, psychosocial counselling) for survivors, including opening hours, contact details, and whether services can be offered remotely, and establish referral linkages. It is important to understand women and girls of marginalised groups and with disabilities are likely to have additional risks and needs. WHO-National Federation of Disabled Nepal (NFDN), Yes We Can Project has set-up district level virtual help desk with woman peer counsellor to address the needs of women and girls with disabilities. The contact numbers are available on request from the National Coordinator of this project (nc@nfdn.org.np).

Remember: Safety, respect, confidentiality and non-discrimination in relation to GBV survivors and those at risk are vital considerations at all times.

3. CONSIDERATIONS FOR MANAGERS

Many women are at the forefront of the COVID-19 response. Study⁶ shows globally, women make up 70 per cent of the health workforce, especially as nurses, midwives and community health volunteers, and account of the majority of service staff in health facilities as cleaners, launderers and caterers. This scenario obtains in Nepal as well. Despite the large number, women are often not reflected in decision-making in response to COVID-19. Further, women are still paid less than their male counterparts and hold fewer leadership positions in the health sector and enjoy lower job security and social protection. Masks and other protective equipment designed and sized for men leave women at greater risk of exposure. The lack of adequate attention to the menstrual hygiene needs of women health workers during long shifts is an added workplace-related challenge.

So, Managers need to be aware of the above gaps and ensure from management aspect if the needs of women especially who are at forefront are prioritized and fulfilled. This means:

- The health care workers and caregivers have access to women/ gender-friendly personal protective equipment (PPE) and menstrual hygiene products, i.e., the different sizes and also the design of the PPE needs to be made available and accessible considering the feminine and menstrual hygiene need.
- Flexible working arrangements need to be made to balance the burden of care specially for pregnant and breastfeeding mothers.
- Women health workers take the leadership and decision-making roles.
- Equal treatment and pay, paid leave and other social protection measures are ensured to women health workers in the public and private sectors.

⁶ UN Women | Explainer: How COVID-19 impacts women and girls

APPENDICES

APPENDIX 1: FIGURES

Appendix 1: Figures

Your 5 Moments for Hand Hygiene





Figure 1: WHO's My 5 Moments for Hand Hygiene

Preventive Measure for COVID-19: Practice Respiratory Hygiene


Make sure you, and the people around you, follow **good respiratory hygiene**.

This means:




Covering your mouth and nose with your bent elbow

or



Covering with tissue when you cough or sneeze


and



Dispose of the used tissue immediately

Why?

Droplets spread virus. By following good respiratory hygiene you protect the people around you from viruses such as cold, flu and COVID-19.







Figure 2: Respiratory Hygiene

58

WASH HANDS WHEN VISIBLY SOILED! OTHERWISE, USE HANDRUB

 **Duration of the entire procedure: 40-60 seconds**



Wet hands with water;



Apply enough soap to cover all hand surfaces;



Rub hands palm to palm;



Right palm over left dorsum with interlaced fingers and vice versa;



Palm to palm with fingers interlaced;



Backs of fingers to opposing palms with fingers interlocked;



Rotational rubbing of left thumb clasped in right palm and vice versa;



Rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa;



Rinse hands with water;



Dry hands thoroughly with a single use towel;



Use towel to turn off faucet;




Your hands are now safe.

Figure 3: How to Handwash?

COVID-19

Coronavirus Disease 2019

HOW TO PUT ON PERSONAL PROTECTIVE EQUIPMENT (PPE)



*Surgical mask or respirator (N95 or similar), depending on the level of care. For aerosol generating procedures (AGP), wear a respirator (N95 or similar)

**e.g. visor, face shield, goggles (consider anti-fog drops or fog-resistant goggles)

PAHO



Pan American
Health
Organization



World Health
Organization
Americas

BE AWARE. PREPARE. ACT.
www.paho.org/coronavirus

Figure 4: Steps to put on PPE (donning)

COVID-19

Coronavirus Disease 2019

HOW TO TAKE OFF PPE

- Avoid contamination of self, others and environment
- Remove the most heavily contaminated items first



Remove gloves
then remove gowns



Perform hand
hygiene



Remove eye or
facial protection
from behind



Remove surgical
mask or respirator
from behind



Perform hand
hygiene

PAHO



Pan American
Health
Organization



World Health
Organization
Member State of the Americas

BE AWARE. PREPARE. ACT.

www.paho.org/coronavirus

Figure 5: Steps to Remove PPE (doffing)




Figure 6. Relaxation Activities for Child and Adolescent (C&A)


FREE HELPLINE NUMBERS FOR MENTAL HEALTH AND PSYCHOSOCIAL SUPPORT			
ORGANIZATION	TOLL FREE NUMBER	ONLINE SERVICE TIME	OTHER HELPLINES
C&A Unit, Kanti Children's Hospital	16600110666	8 am to 5 pm	<div><div>Kathmandu University School of Medical Sciences, Dhulikhel Hospital (Psychiatric Consultation)</div><div>9851310998 9801315214 (Office Hour)</div></div> <div><div>TU Teaching Hospital (Psychological Service Hotline)</div><div>9849925963 (Office Hour)</div></div> <div><div>Unity in Health Nepal (Psychological Support for Frontline Health Workers & Public)</div><div>9843643433 9801166258 9881365991 (Office Hour)</div></div> <div><div>Psychbigyaan Network - PNN (Psychological Support for Organizations & Public)</div><div>9860937293 (Office Hour)</div></div> <div><div>Chhahari Nepal for Mental Health (Psychosocial Support for Street People)</div><div>9851183773, 9848068221 (Office Hour)</div></div> <div><div> Government of Nepal Ministry of Health and Population</div></div>
CMC-Nepal	16600185080	9 am - 5 pm	
Health Foundation Nepal	16608256003	8 am to 10 pm	
KOPILA-Nepal	16606152005	8 am to 8 pm (Open on Saturday)	
KOSHISH	16600122322	Open on Saturday	
Kashish Nepal (Gandaki)	16606152007	Open on Saturday	
National Women Commission	1145	24 hours (Open on Saturday)	
Nepal Red Cross Society	1130	7 am - 7 pm	
Patan Mental Hospital (Suicide Prevention Hotline)	1166	24 hours (Open on Saturday)	
TPO Nepal	16600102005	8 am - 6 pm (Open on Saturday)	
TU Teaching Hospital (Suicide Prevention Hotline)	16600121600	24 hours (Open on Saturday)	

Figure 7. Service Provider Related Information



Fig 8. Communication Board for COVID-19 Facilities




Fig 9. Calm Down Clock





Fig 10. Calm Down Dice Web

APPENDIX 2: FLOW CHARTS


Flow chart 1:

Approach to baby born to suspected/RT-PCR positive/COVID antigen positive mothers




Flow chart 2:

Approach to a symptomatic baby born to suspected/RT-PCR positive/COVID antigen positive mothers



Flow chart 3:

Discontinuing isolation and precautions guidelines for suspected or confirmed COVID mothers




Recommendations for breastfeeding and breast milk feeds in the context of COVID-19

- Breastfeeding should be encouraged and continued. If direct breastfeeding is not possible, expressed breast milk should be given to baby unless contraindicated
- Mother should wash their hands using soap and water before touching their child or expressing breast milk. If soap and water are not available, use hand sanitizer with at least 60% alcohol.
- Mother should wear a mask when they are less than 6 feet from the child (including when feeding at the breast or feeding from a bottle) and when expressing breast milk
- If mother is very sick to breastfeed or express breast milk, donor human milk, wet nursing or appropriate breast milk substitutes should be considered based on feasibility, safety, sustainability, cultural context, acceptability to mother and service availability.

Flow Chart 4:

Diagnostic algorithm for the evaluation of patient with suspected MIS-C



First line investigations: CBC, LFT, RFT, PT INR, aPTT, Blood gas analysis, Blood glucose, CRP and or ESR, SARS-Co-2 serology, RT-PCR. Consider and send investigations for other infections: Malaria, Dengue, Blood C/S, Scrub typhus, Leptospirosis, etc.


Positive first line investigations: CRP>5mg/dl and or ESR >40 mm/hr PLUS ALC <1000/ μ L, or Platelets<150,000/ μ L, or Na<135mEq/L, or Neutrophilia, or Hypoalbuminemia.

Second line investigations: Cardiac (ECG, ECHO, BNP, Troponin T), inflammatory markers (Procalcitonin, D-dimer, Fibrinogen, LDH, Triglyceride, IL-6, Ferritin), Peripheral blood smear.

Note: Investigations should be sent as per its availability/feasibility at the treating center.

Consider and send investigations for other infections: Malaria, Dengue, Blood C/S, Scrub typhus, Leptospirosis
Positive first line investigations: CRP>5mg/dl and or ESR >40 mm/hr PLUS ALC <1000/cu mm or Platelets<150000/cu mm, or Na<135mEq/L, or Neutrophilia, or Hypoalbuminemia

Flow Chart 5:
Algorithm for the management of MIS-C



APPENDIX 3: MULTISYSTEM INFLAMMATORY SYNDROME IN CHILDREN (MIS-C)

I. Clinical manifestations of MIS-C

SYMPTOMS	SIGNS
<ul style="list-style-type: none"> • Fever • Gastrointestinal: abdominal pain, vomiting, diarrhea • Rash • Conjunctivitis • Mucous membrane involvement • Neurological: headache, lethargy, confusion • Respiratory: tachypnea, labored breathing • Sore throat • Myalgias • Swollen hands/feet • Lymphadenopathy 	<ul style="list-style-type: none"> • Shock • Features of Kawasaki disease • Myocardial dysfunction (by echo cardiogram or elevated troponin/BNP) • Arrhythmia • Acute respiratory failure requiring noninvasive or invasive ventilation • Acute kidney injury • Serositis (small pleural, pericardial, and ascitic effusions) • Hepatitis or hepatomegaly • Encephalopathy, seizures, coma, or meningo-encephalitis
Laboratory Investigations Abnormal blood cell counts <ul style="list-style-type: none"> • Lymphocytopenia • Neutrophilia • Mild anemia • Thrombocytopenia • Elevated inflammatory markers • C-reactive protein • Erythrocyte sedimentation rate • D-dimer (if available) 	Radiology Echocardiogram <ul style="list-style-type: none"> • Depressed LV function • Coronary artery dilation/aneurysm • Other findings can include mitral regurgitation and pericardial effusion

<ul style="list-style-type: none"> • Erythrocyte sedimentation rate • D-dimer (if available) • Fibrinogen (if available) • Ferritin • Procalcitonin (if available) <p>Elevated cardiac markers</p> <ul style="list-style-type: none"> • Troponin • BNP (if available) • Hypoalbuminemia • Mildly elevated liver enzymes • Elevated lactate dehydrogenase • Hypertriglyceridemia 	<p>Chest radiograph</p> <ul style="list-style-type: none"> • Normal in many patients • Abnormal findings: small pleural effusions, patchy consolidations, focal consolidation and atelectasis <p>Chest CT</p> <ul style="list-style-type: none"> • Findings generally similar to those on chest radiograph • A few patients had nodular ground-glass opacification <p>Abdominal imaging (ultrasound and/or CT)</p> <ul style="list-style-type: none"> • Ascites • Bowel and mesenteric inflammation • Terminal ileitis • Mesenteric adenopathy/adenitis • Pericholecystic edema
---	---

II. Distinction features between MIS-C and Kawasaki disease (KD)

	MIS-C	KD
Age	Older children & adolescents	Infants & young children
Ethnicity	Increased incidence in African, Afro-Caribbean and Hispanic descent	Common in East Asian children
Gastrointestinal symptoms	More common	Less prominent
Shock	More common	Less common
Myocardial dysfunction (arrhythmias & ventricular dysfunction)	More common	Less common

Neurological symptoms	More common	Less common
Inflammatory markers	Markedly elevated	Elevated
Platelet count	Thrombocytopenia	Thrombocytosis
Coronary artery involvement	Risk present	Risk present

III. Definition of organ system involvement

Gastrointestinal: nausea/vomiting, diarrhea, abdominal pain, appendicitis, pancreatitis, hepatitis, gallbladder hydrops or edema
Cardiovascular: hypotension or shock, cardiac dysrhythmia or arrhythmia, ejection fraction <55%, pulmonary edema due to left heart failure, coronary artery z score ≥ 2.5 , pericarditis or pericardial effusion or valvulitis, B-type natriuretic peptide (BNP) >400 pg/mL, elevated troponin, receipt of vasopressor or vasoactive support, receipt of cardiopulmonary resuscitation (CPR)
Hematologic: Total white blood cell <4,000, anemia for age, platelet count <150,000 / μ L, deep vein thrombosis, pulmonary embolism, hemolysis, bleeding or prolonged PT/aPTT, ischemia of an extremity
Mucocutaneous: bilateral conjunctival injection, oral mucosal changes, rash or skin ulcers, 'COVID' toes, swollen red cracked lips, erythema of palms or soles, edema of hands or feet, periungual (nails) desquamation
Respiratory: receipt of mechanical ventilation or any type of supplemental oxygen (or increased support for patients receiving respiratory support at baseline), severe bronchospasm requiring continuous bronchodilators or pulmonary infiltrates on chest radiograph, lower respiratory infection, pleural effusion, pneumothorax or other signs of barotrauma, pulmonary hemorrhage, chest-tube or drainage required
Musculoskeletal 23% (more frequent in teens): arthritis or arthralgia, myositis or myalgia
Renal: acute kidney injury with or without dialysis
Neurologic: stroke or acute intracranial hemorrhage, seizures, encephalitis, aseptic meningitis, or demyelinating disorder, altered mental status, suspected meningitis with negative culture

IV. Differentiating features between MIS-C and Severe COVID in children

Features	MIS C	Severe COVID-19
Age group (years)	6-12	0-5
Severe pulmonary involvement	Less common	Common
Multiple Organ Dysfunction Syndrome (MODS) <ul style="list-style-type: none"> • Cardiovascular • Gastrointestinal • Mucocutaneous 	Common	Less common
<ul style="list-style-type: none"> • Laboratory parameters • Thrombocytopenia ($150 \times 10^3/\mu\text{L}$) • Elevated CRP ($>100\text{mg/dl}$) • Hypoalbuminemia • Neutrophil lymphocyte ratio (>5) 	Common	Less common

REFERENCES

1. Infection prevention and control of epidemic- and pandemic-prone acute respiratory infections in health care: WHO guidelines. Geneva: World Health Organization; 2014 (available at: https://apps.who.int/iris/bitstream/handle/10665/112656/9789241507134_eng.pdf, accessed 14 Feb 2021).
2. Infection prevention and control during health care when coronavirus disease (COVID-19) is suspected or confirmed: Interim guidance: WHO. Geneva: World Health Organization; 12 July 2021 (available at: <https://www.who.int/publications/i/item/WHO-2019-nCoV-IPC-2021.1>, accessed on 11 Nov 2021).
3. WHO guidelines on hand hygiene in health care: first global patient safety challenge - clean care is safer care. Geneva: World Health Organization; 2009 (available at: https://www.who.int/gpsc/5may/tools/who_guidelines-handhygiene_summary.pdf, accessed 20 Feb 2021).
4. Hand Hygiene: Why, How & When? Geneva: World Health Organization; 2009 (available at: https://www.who.int/gpsc/5may/Hand_Hygiene_Why_How_and_When_Brochure.pdf, accessed on 13 Feb 2021).
5. Advice on the use of masks for children in the community in the context of COVID-19. Annex to the Advice on the use of masks in the context of COVID-19: World Health Organization; 21 Aug 2020 (available at: https://www.who.int/publications/i/item/WHO-2019-nCoV-IPC_Masks-Children-2020.1).
6. How to put on and take off personal protective equipment (PPE). Geneva: World Health Organization; 2020 (available at: <https://www.who.int/csr/resources/publications/putontake-offPPE/en/>, accessed 15 Feb 2021).
7. Rational use of personal protective equipment for coronavirus disease 2019 (COVID-19). Geneva: World Health Organization; 2020 (available at: [https://www.who.int/publications/i/item/rational-use-of-personal-protective-equipment-for-coronavirus-disease-\(covid-19\)-and-considerations-during-severe-shortages](https://www.who.int/publications/i/item/rational-use-of-personal-protective-equipment-for-coronavirus-disease-(covid-19)-and-considerations-during-severe-shortages), accessed 21 Feb 2021).
8. Honda H, Iwata K. Personal protective equipment and improving compliance among health-care workers in high-risk settings. *Curr Opin Infect Dis.* 2016;29(4):400-406.
9. Guidelines on core components of infection prevention and control programmes at the national and acute health care facility level. Geneva: World Health Organization; 2016 (available at: <https://www.who.int/gpsc/ipc-components/en/>, accessed 14 Feb 2021).
10. CDC and ICAN. Best Practices for Environmental Cleaning in Healthcare Facilities in Resource-Limited Settings. Atlanta, GA: US Department of Health and Human Services, CDC; Cape Town, South Africa: Infection Control Africa Network; 2019 (available at: <https://www.cdc.gov/hai/pdfs/resource-limited/environmental-cleaning-RLS-H.pdf>, accessed 14 Feb 2021).
11. Cleaning and disinfection of environmental surfaces in the context of COVID-19. Geneva: World Health Organization; 2020 (available at: <https://www.who.int/publications/i/item/cleaning-and-disinfection-of-environmental-surfaces-in-the-context-of-covid-19>, accessed 21 Feb 2021).
12. Rutala, WA, Weber, DJ., 2019. Best practices for disinfection of noncritical environmental surfaces and equipment in health care facilities: A bundle approach. *Am J Infect Control* 47, A96-A105.
13. Decontamination and reprocessing of medical devices for health care facilities. Geneva: World Health Organization; 2016 (available at: <https://www.who.int/infection-prevention/publications/decontamination/en/>, accessed 16 Feb 2021).
14. Water, sanitation, hygiene, and waste management for the COVID-19 virus: interim guidance. Geneva: World Health Organization; 2020 (available at: <https://apps.who.int/iris/handle/10665/331499>, accessed 16 Feb 2021).

15. Modes of transmission of virus causing COVID-19: implications for IPC precaution recommendations 29 March 2020 [Internet]. Geneva: World Health Organization; 2020 (available at: <https://www.who.int/news-room/commentaries/detail/modes-of-transmission-of-virus-causing-covid-19-implications-for-ipc-precaution-recommendations>, assessed 21 Feb 2021).
16. Atkinson J, Chartier Y, Pessoa-Silva CK, Jensen P, Li Y, Seto WH, editors. Natural ventilation for infection control in health care settings. Geneva: World Health Organization; 2009 (available at: https://www.who.int/water_sanitation_health/publications/natural_ventilation.pdf, accessed 16 Feb 2021).
17. Use of chest imaging in COVID-19. Geneva: World Health Organization; 2020 (available at: <https://www.who.int/publications/i/item/use-of-chest-imaging-in-covid-19>, accessed 16 Feb 2021).
18. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ, COVID-19 Systematic Urgent Review Group Effort (SURGE) study authors. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet*. 2020 June 1. doi: 10.1016/S0140-6736(20)31142-9/
19. Advice on the use of masks in the context of COVID-19. Geneva: World Health Organization; 2020 (available at: <https://apps.who.int/iris/handle/10665/331693>, accessed 16 Feb 2021).
20. Clinical management of COVID-19. Geneva: World Health Organization; 2020 (available at: <https://apps.who.int/iris/handle/10665/332196>, accessed on 15 Feb 2021).
21. Lan L, Xu D, Ye G, Xia C, Wang S, Li Y, Xu H. Positive RT-PCR Test Results in Patients Recovered From COVID-19. *JAMA*. 2020 Apr 21;323(15):1502-3.
22. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *CADTH Technol Overv*. 2013;3(1):e3101.
23. Hui DS. Epidemic and emerging coronaviruses (severe acute respiratory syndrome and Middle East respiratory syndrome). *Clin Chest Med*. 2017 Mar;38(1):71-86. doi: 10.1016/j.ccm.2016.11.007.
24. Heinzerling A, Stuckey MJ, Scheuer T, et al. Transmission of COVID-19 to health care personnel during exposures to a hospitalized patient - Solano County, California, February 2020. *MMWR Morb Mortal Wkly Rep* 2020 Apr 17;69(15):472-6. doi: <http://dx.doi.org/10.15585/mmwr.mm6915e5>.
25. How to perform a particulate respirator seal check? Geneva: World Health Organization; 2008 (available at: <http://www.who.int/csr/resources/publications/respiratorsealcheck/en/>, accessed 16 Feb 2021).
26. Laboratory biosafety guidance related to coronavirus disease (COVID-19). Geneva: World Health Organization; 2020 [available at: [https://www.who.int/publications/i/item/laboratory-bio-safety-guidance-related-to-coronavirus-disease-\(COVID-19\)](https://www.who.int/publications/i/item/laboratory-bio-safety-guidance-related-to-coronavirus-disease-(COVID-19)), accessed 16 Feb 2021].
27. The Pandemic Surgery Guidance Consortium (PSGC). COVID-19: Pandemic surgery guidance. EDP Sciences, 2020 (available at: https://www.4open-sciences.org/articles/fopen/full_html/2020/01/fopen200002s/fopen200002s.html, accessed 21 Feb 2021).
28. COVIDSurg Collaborative. Mortality and pulmonary complications in patients undergoing surgery with perioperative SARS-CoV-2 infection: an international cohort study. *Lancet*. 2020 May 29;396(10243):27-38. doi: 10.1016/S0140-6736(20)31182-X.
29. Judson SD, Munster VJ. Nosocomial Transmission of Emerging Viruses via Aerosol-Generating Medical Procedures. *Viruses*. 2019 Oct;11(10):940. Published 2019 Oct 12. doi:10.3390/v11100940.
30. Community-based health care, including outreach and campaigns, in the context of the COVID-19 pandemic. WHO and UNICEF, 2020 (available at: <https://www.unicef.org/documents/community-based-health-care-outreach-campaigns-covid-19-pandemic>, accessed 16 Feb 2021).
31. Telemedicine opportunities and development in member states. Geneva: World Health Organization; 2010 (available at https://www.who.int/goe/publications/goe_telemedicine_2010.pdf, accessed 18 Feb 2021).
32. Infection prevention and control for the safe management of a dead body in the context of COVID-19: interim guidance. Geneva: World Health Organization; 2020 (available at: <https://>

www.who.int/publications/i/item/infection-prevention-and-control-for-the-safe-management-of-a-dead-body-in-the-context-of-covid-19-interim-guidance, accessed 18 Feb 2021).

33. World Health Organization. Director-General's remarks at the media briefing on 2019-nCoV on 11 February 2020. <http://www.who.int/dg/speeches/detail/who-director-general-s-remarks-at-the-media-briefing-on-2019-ncov-on-11-february-2020>

34. World Health Organization. COVID-19 clinical management. Living guidance. January 25, 2021. Available at: <https://www.who.int/publications/i/item/WHO-2019-nCoV-clinical-2021-1>

35. World Health Organization. Clinical management of severe acute respiratory infection (SARI) when COVID-19 disease is suspected. Interim guidance 13 March 2020

36. Centers for Disease Control and Prevention. Information for healthcare professionals about coronavirus (COVID-19). <https://www.cdc.gov/coronavirus/2019-nCoV/hcp/index.html>

37. Care for Breastfeeding People Interim Guidance on Breastfeeding and Breast Milk Feeds in the Context of COVID-19. Updated Dec. 3, 2020.

38. South AM, Brady TM, Flynn JT. ACE2 (Angiotensin-Converting Enzyme 2), COVID-19, and ACE Inhibitor and Ang II (Angiotensin II) Receptor Blocker Use During the Pandemic: The Pediatric Perspective. *Hypertension* 2020; 76:16.

39. Koeckerling D, et al. *Thorax* October 2020 Vol 75 No 10. *CLINICS* 2020;75:e1894

40. Dyah KW and Arya KM. Management of children with COVID-19. *CEP* Vol. 63, No. 9, 345-354, 2020, <https://doi.org/10.3345/cep.2020.00913>

41. Carlotti APCP et al. Management Pediatric COVID-19.in

42. S Kache et al. COVID-19 PICU guidelines: for high- and limited-resource settings. *Pediatric Research* (2020) 88:705-716; <https://doi.org/10.1038/s41390-020-1053-9>

43. www.askgileadmedical.com

44. Marini JJ, Gattinoni L. Management of COVID-19 Respiratory Distress. *JAMA* June 9, 2020 Volume 323, Number 22. doi:10.1001/jama.2020.6825.

45. Mittermaier M et al. Evaluation of PEEP and prone positioning in early COVID-19 ARDS. *EClinicalMedicine* 28 (2020) 100579. <https://doi.org/10.1016/j.eclinm.2020.100579>.

46. Sankar J, Dhochak N, Kabra SK and Lodha R. *Indian J Pediatr* (442 June 2020) 87(6):433-442

47. Loi M et al. COVID-19 anticoagulation recommendations in children. *Pediatric blood and cancer*. Volume 67, Issue 9, September 2020 e28485. <https://doi.org/10.1002/pbc.28485>

48. Thachil J, Tang N, Gando S, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost*. 2020; 18(5): 1023- 1026.

49. Centers for Disease Control and Prevention. Interim Guidance for Antigen Testing for SARS-CoV-2. Updated Dec. 16, 2020 <https://www.cdc.gov/coronavirus/2019-ncov/lab/resources/antigen-tests-guidelines.html>.

50. Centers for Disease Control and Prevention. When You Can be Around Others After You Had or Likely Had COVID-19. Updated Feb. 11, 2021.

51. American Academy of Pediatrics. Critical Update on COVID19/COVID-19 Interim Guidance/ Frequently Asked Questions: Interfacility Transport of the critically ill neonatal or pediatric patient with suspected or confirmed COVID 19.

52. MANU SUNDARAM, NAMITA RAVIKUMAR, ARUN BANSAL, KARTHI NALLASAMY, BASAVARAJA GV, RAKESH LODHA et al, Novel Coronavirus 2019 (2019-nCoV) Infection: Part II - Respiratory-Support in the Pediatric Intensive Care Unit in Resource-limited Settings, *Indian Pediatrics*, VOLUME 57, APRIL 15, 2020

53. Ira M Cheifetz. Pediatric ARDS. *Respiratory Care* June 2017, 62 (6) 718-731; DOI: <https://doi.org/10.4187/respcare.05591>

54. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. Available at <https://www.covid19treatmentguidelines.nih.gov/>. Accessed on 10 July 2021

55. Guidelines for Management of COVID-19 in Children (below 18 years) .Ministry of Health & Family Welfare Government of India, 18th June 2021.

56. Centers for Disease Control and Prevention's (CDC) Evaluation and Management Considerations for Neonates at Risk for COVID-19.

- 57.WHO COVID-19 Clinical management. Feeding and caring for infants and young children of mothers with COVID-19. Living guidance. 25 January 2021.
- 58.Walker KF, O'Donoghue K, Grace N, Dorling J, Comeau JL, Li W, Thornton JG Maternal transmission of SARS-CoV-2 to the neonate, and possible routes for such transmission: a systematic review and critical analysis. BJOG. 2020;127(11):1324. Epub 2020 Jul 22.Cojocaru L, Crimmins S, Sundararajan S, Goetzinger K, Elsamadicy E, Lankford A, Turan OM, Turan S, An initiative to evaluate the safety of maternal bonding in patients with SARS-CoV-2 infection. J Matern Fetal Neonatal Med. 2020.
- 59.Gale C, Quigley MA, Placzek A, Knight M, Ladhani S, Draper ES, Sharkey D, Doherty C, Mactier H, Kurinczuk JJ. Characteristics and outcomes of neonatal SARS-CoV-2 infection in the UK: a prospective national cohort study using active surveillance. Lancet Child Adolesc Health. 2021 Feb;5(2):113-121. doi: 10.1016/S2352-4642(20)30342-4. Epub 2020 Nov 9.
- 60.Lana A. Shaiba, Adnan Hadid, Khalid A. Altirkawi, Hind M. Bakheet, Aminah Mohammed Alherz et al. Case Report: Neonatal Multisystem Inflammatory Syndrome Associated With SARS-CoV-2 Exposure in Two Cases From Saudi Arabia. Front. Pediatr., 13 May 2021 | <https://doi.org/10.3389/fped.2021.652857>
- 61.Henderson LA, Canna SW, Friedman KG, Gorelik M, Lapidus SK, Bassiri H, et al. American College of Rheumatology Clinical Guidance for Pediatric Patients with Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with SARS-CoV2 and Hyperinflammation in COVID-19. Version 1. Arthritis Rheumatol 2020; 72: 1791-1805. doi: <https://onlinelibrary.wiley.com/doi/10.1002/art.41454>.
- 62.Royal College of Paediatrics and Child Health (RCPCH) Guidance: Paediatric multisystem inflammatory syndrome temporally associated with COVID19 <https://www.rcpch.ac.uk/sites/default/files/2020-05/COVID-19-Paediatric-multisystem-%20inflammatory%20syndrome-20200501.pdf>
- 63.Centres for Disease Control and Prevention (CDC): Information for Healthcare Providers about Multisystem Inflammatory Syndrome in Children (MIS-C). <https://www.cdc.gov/mis-c/index.html>
- 64.Word Health Organization (WHO): Multisystem inflammatory syndrome in children and adolescents with COVID-19: Scientific brief 15 May 2020 <https://www.who.int/publications-detail-redirect/multisystem-inflammatory-syndrome-in-children-and-adolescents-with-covid-19>
- 65.Feldstein LR, Rose EB, Horwitz SM, Collins JP, Newhams MM, Son MBF, Newburger JW, Kleinman LC et al; Overcoming COVID-19 Investigators; CDC COVID-19 Response Team. Multisystem Inflammatory Syndrome in U.S. Children and Adolescents. N Engl J Med. 2020 Jul 23;383(4):334-346. doi: 10.1056/NEJMoa2021680. Epub 2020 Jun 29. PMID: 32598831; PMCID: PMC7346765
- 66.Dufort EM, Koumans EH, Chow EJ, Rosenthal EM, Muse A, Rowlands J, Barranco MA et al. Multisystem Inflammatory Syndrome in Children in New York State. N Engl J Med. 2020;383(4):347. PMID 32598830
- 67.Davies P, Evans C, Kanthimathinathan HK, Lillie J, Brierley J, Waters G, Johnson M, Griffiths B et al. Intensive care admissions of children with paediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 (PIMS-TS) in the UK: a multicentre observational study. Lancet Child Adolesc Health. 2020;4(9):669. PMID 32653054
- 68.American College of Rheumatology Clinical Guidance for Pediatric Patients with Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with SARS-CoV-2 and Hyperinflammation in COVID-19. Version 2. Arthritis Rheumatol. <https://doi.org/10.1002/art.41616>
- 69.COVID-19 pathway v4. COVID19Pathway@seattlechildrens.org
- 70.Clinical Updates on COVID-19/ COVID-19 interim guidance/ Multisystem Inflammatory Syndrome in Children (MIS-C) Interim Guidance. Last updated 11/17/2020. American Academy of Pediatrics.
- 71.Jiang L, Tang K, Levin M, Irfan O, Morris SK, Wilson K, et al. COVID-19 and multisystem inflammatory syndrome in children and adolescents. Lancet Infect Dis. 2020; 20:276-88.
- 72.Best available treatment study for inflammatory conditions associated with COVID-19. Available from: <http://www.isrctn.com/ISRCTN69546370>
- 73.McMurray JC, May JW, Cunningham MW, Jones OY. Multisystem Inflammatory Syndrome

Management protocol of Children with COVID-19

in Children (MIS-C), a Post-viral Myocarditis and Systemic Vasculitis—A Critical Review of Its Pathogenesis and Treatment. *Front Pediatr*. 2020 16; 8:626182.

74.Guideline for the management of Paediatric multisystem inflammatory syndrome temporally associated with COVID-19. 2020;10.

75.Hennon TR, Penque MD, Abdul-Aziz R, Alibrahim OS, McGreevy MB, Prout AJ, et al. COVID-19 associated Multisystem Inflammatory Syndrome in Children (MIS-C) guidelines; a Western New York approach. *ProgPediatrCardiol*. 2020; 57:101232.

76.Ouldali N, Toubiana J, Antona D, Javouhey E, Madhi F, Lorrot M, et al. Association of Intravenous Immunoglobulins Plus Methylprednisolone vs Immunoglobulins Alone With Course of Fever in Multisystem Inflammatory Syndrome in Children. *JAMA*. 2021

77.Interim Management protocol for COVID-19 in children. AIIMS, New Delhi;14 April 2021.

78.Multisystem Inflammatory Syndrome in Children (MIS-C): Statement by Indian Academy of Pediatrics; April 2021.

79.Reference: Feldstein LR, Tenforde MW, Friedman KG, et al. Characteristics and Outcomes of US Children and Adolescents with Multisystem Inflammatory Syndrome in Children (MIS-C) Compared with Severe Acute COVID-19. *JAMA*.2021; 325:1074–1087. doi:10.1001/jama.2021.2091

80.Central Bureau of Statistics of Nepal. Statistical Yearbook Nepal 2015 [Internet]. 2016. Available from: <http://cbs.gov.np/image/data/2017/Statistical Year Book 2015.pdf>

81.Pradhan TR. Nepal goes under lockdown for a week starting 6am Tuesday. The Kathmandu Post [Internet]. 2020 Mar 23 [cited 2020 Nov 13]; Available from: <https://kathmandupost.com/national/2020/03/23/nepal-goes-under-lockdown-for-a-week-starting-6am-tuesday>

82.Sharma V, Reina Ortiz M, Sharma N. Risk and Protective Factors for Adolescent and Young Adult Mental Health within the Context of COVID-19: A Perspective From Nepal [Internet]. Vol. 67, *Journal of Adolescent Health*. Elsevier USA; 2020 [cited 2020 Nov 13]. p. 135–7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7237905/>

83.Dhonju, G., Kunwar, A. R., Karki, U., Devkota, N., Bista, I., &Sah, R. (2021). Identification and management of CoViD-19 related child and adolescent mental health problems: a multi-tier intervention model. *Frontiers in public health*, 8, 992.

84.Reference: Clinical guide for front line staff to support the management of patients with a learning disability, autism or both during the COVID-19 pandemic – relevant to all clinical specialities, NHS, November 2020, revised May 2021

LIST OF CONTRIBUTORS

Dr. Pawanjung Rayamajhi,
Director, CSD

Ms. Roshani Laxmi Tuitui,
Director, NSSD

Dr. Narendra Kumar Khanal,
Senior Consultant Medical Generalist, CSD

Dr. Prakash Budhathoki,
Senior Consultant Dental Surgeon, CSD

Dr. Pomawati Thapa,
Senior Consultant Medical Generalist, CSD

Dr. Lila Bahadur Basnet,
Community Physician, CSD

Mr. Bharat Mani Marhatta,
Senior Pharmacy Officer, CSD

Mr. Pradip Kumar Yadav,
Public Health Officer, CSD

Ms. Shirley McQuen Patterson
WHO

Prof. Dr. Shital Adhikari,
WHO

Dr. Subash Neupane,
WHO

Dr. Sudesha Khadka,
WHO

Mr. Kamaraj Devapitchai
WHO

Ms. Sadhana Paudel
WHO

Ms. Chahana Singh,
UNICEF

Dr. Samikshya Neupane Dulal,
UNICEF

Mr. Parashuram Shrestha,
UNICEF

NOTES

[illegible]

Supported by:



Nepal Paediatric Society
Dedicated to the health of allChildren



**World Health
Organization**
Nepal

unicef 
for every child